



Tumour Review



PREDICTIVE *BRCA* GENETIC TESTING IN ITALIAN PATIENTS WITH BREAST CANCER: A POSITION PAPER OF ITALIAN SCIENTIFIC SOCIETIES [Italian Association of Medical Oncology(AIOM); Italian Association of Radiotherapy and Clinical Oncology (AIRO); Italian National Association of Breast Surgeons (ANISC); Italian Society of Pathological Anatomy and Diagnostic Cytology (SIAPeC-IAP); Italian Society of Surgical Oncology (SICO); Italian Society of Human Genetic (SIGU); Italian Society of General Practice (SIMG); Italian Society of Medical and Interventional Radiology (SIRM)]

L. Cortesi^{a,b,*}, F. Miglietta^c, L. Arecco^d, D. Bernardi^{e,f}, L. Biganzoli^g, L. Del Mastro^{d,h}, MV. Dieci^c, J. Fogliettaⁱ, L. Fortunato^j, P. Franco^{k,l}, E. Lucci Cordisco^m, P. Mantellini^{n,o}, C. Marchio^{p,q}, B. Meduri^r, G. Micallo^s, A. Musolino^t, A. Salvetti^u, D. Turchetti^v, A. Zambelli^{w,x}, C. Angiolini^y, S. Cinieri^z, PF. Conte^{aa}, G. Curigliano^{ab,ac}, M. Dessena^{ad}, A. Fabi^{ae}, MC. Leonardi^{af}, I. Meattini^{ag,ah}, S. Montemezzi^{ai}, F. Puglisi^{aj,ak}, A. Sapino^{p,q}, M. Taffurelli^{al}, M. Cinquini^{am}, S. Gori^{an}

^a Medical Oncology, Azienda USL-IRCCS di Reggio Emilia, Reggio Emilia, Italy

^b Department of Medical and Surgical Sciences, University of Modena and Reggio Emilia, 41124 Modena, Italy

^c Oncologia Medica 2, IRCCS Istituto Oncologico Veneto, DiSCOG Università degli Studi di Padova, Padova, Italy

^d U.O.C. Clinica di Oncologia Medica, IRCCS Ospedale Policlinico San Martino, Genova, Italy

^e Department of Biomedical Sciences, Humanitas University, Via Rita Levi Montalcini 4, 20090 Pieve Emanuele, Milan, Italy

^f IRCCS Humanitas Research Hospital, Via Manzoni 56, 20089 Rozzano, Milan, Italy

^g Department of Oncology, Hospital of Prato, Azienda USL Toscana Centro, Prato, Italy

^h Department of Internal Medicine and Medical Specialties, Università di Genova, Genova, Italy

ⁱ Oncologia Medica e Trasfusione, Ospedale Santa Maria di Terni, Italy

^j Breast Centre, Azienda Ospedaliera San Giovanni-Addolorata, 00184 Rome, Italy

^k Department of Translational Medicine (DIMET), University of Eastern Piedmont, Novara, Italy

^l Department of Radiation Oncology, "Maggiore della Carità" University Hospital, Novara, Italy

^m Fondazione Policlinico Universitario A. Gemelli IRCCS, UOC Genetica Medica, Rome, Italy

ⁿ Sezione di Medicina Genomica, Dipartimento di Scienze della Vita e Sanità Pubblica, Università Cattolica del Sacro Cuore, Rome, Italy

^o Institute for Cancer Research, Prevention and Clinical Network (ISPRO), Florence, Italy

^p Department of Medical Sciences, University of Turin, Turin, Italy

^q Candiolo Cancer Institute, FPO-IRCCS, Candiolo, Italy

^r Department of Radiation Oncology, University Hospital of Modena, Modena, Italy

^s Hematology-Oncology and Stem Cell Transplantation Unit, Istituto Nazionale Tumori, Fondazione G. Pascale, IRCCS, Naples, Italy

^t Medical Oncology, Breast & GYN Unit, IRCCS Istituto Romagnolo per lo Studio dei Tumori (IRST) "Dino Amadori", Meldola, Italy

^u General Practitioner, ASL 9 Grosseto, Italy

^v Medical Genetics Unit, IRCCS Azienda Ospedaliero-Universitaria di Bologna, Bologna, Italy

^w Medical Oncology, Ospedale Papa Giovanni XXIII, Bergamo, Italy

^x Department of Medicine and Surgery, University of Milano Bicocca, Italy

^y Breast Medical Oncology, Oncology Department, Careggi Hospital, Firenze, Italy

^z UOC Oncologia medica e Breast unit Ospedale Perrino, Brindisi, Fondazione AIOM, Italy

* Corresponding author at: Medical Oncology, Azienda USL-IRCCS di Reggio Emilia, 42123 Reggio Emilia Italy; and University of Modena and Reggio Emilia, Via Università 41121, Modena, Italy.

E-mail address: hbc@unimore.it (L. Cortesi).

<https://doi.org/10.1016/j.ctrv.2025.102976>

Received 10 March 2025; Received in revised form 9 May 2025; Accepted 8 June 2025

Available online 15 June 2025

0305-7372/© 2025 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

^{aa} IRCCS San Camillo-Venice Lido, Italy^{ab} Istituto Europeo di Oncologia, IRCCS, Milano, Italy^{ac} Dipartimento di Oncologia ed Emato-Oncologia, Università di Milano, Milano, Italy^{ad} S.S. Senologia Chirurgica, Chirurgia Polispécialistica, Policlinico Universitario di Monserrato, Azienda Ospedaliera Universitaria, Cagliari, Italy^{ae} Precision Medicine Unit in Senology, Fondazione Policlinico Universitario Agostino Gemelli IRCCS, Largo Agostino Gemelli, 8, 00168 Roma, Italy^{af} Division of Radiation Oncology, IEO European Institute of Oncology, IRCCS, Milan, Italy^{ag} Department of Experimental and Clinical Biomedical Sciences “M. Serio” - University of Florence, Italy^{ah} Breast Unit & Radiation Oncology Unit, Azienda Ospedaliero-Universitaria Careggi, Florence, Italy^{ai} Department of Radiology, IRCCS Sacro Cuore Don Calabria Hospital, Negrar di Valpolicella, Verona, Italy^{aj} CRO Aviano, National Cancer Institute, IRCCS, Aviano, Italy^{ak} Department of Medicine, University of Udine, Udine, Italy^{al} Alma Mater Studiorum-Università di Bologna, Italy^{am} Laboratorio di metodologia delle revisioni sistematiche e produzione di linee guida, Istituto di Ricerche Farmacologiche Mario Negri IRCCS, Milano, Italy^{an} Medical Oncology, IRCCS Sacro Cuore Don Calabria Hospital, Negrar di Valpolicella, Verona, Italy

ARTICLE INFO

Keywords:

BRCA1/2

Germline pathogenic variants (gPV)

PARP inhibitors (PARPi)

Mainstreaming cancer genetics (MCG)

ABSTRACT

The introduction of Poly (ADP-ribose) Polymerase (PARP) inhibitors in both metastatic and early-stage breast cancer (BC) treatment has led to the emergence of Mainstreaming Cancer Genetics (MCG) as a new approach to genetic counselling, predictive of therapy outcomes. Therefore, the *BRCA* testing criteria for therapeutic purposes require further implementation. This position paper outlines the Italian indications for predictive genetic testing, approved by a multidisciplinary Expert Panel representing major scientific societies involved in BC treatment in Italy.

We utilized the Grading of Recommendations, Assessment, Development and Evaluation (GRADE) approach, and framed clinical questions as population, intervention, comparison and outcome (PICO). The final recommendations were determined through a voting process, covering key topics such as eligibility criteria for onco-genetic counselling, the role of contralateral prophylactic mastectomy (CPM) in patients harboring *BRCA1/2* germline pathogenic variants (gPV), and the positioning of predictive *BRCA1/2* test.

As results, the Expert Panel defined three distinct patient groups eligible for onco-genetic counselling, based on the *a priori* likelihood of carrying a gPV and the purpose of testing (predictive vs preventive). A conditional recommendation in favor of CPM in patients with a history of surgically treated BC and a *BRCA* gPV was suggested. Finally, a multidisciplinary pathway for *BRCA* testing was proposed, for patients with triple negative and hormone receptor-positive (HR+)/HER2-negative (HER2-) BC.

In conclusion, the predictive *BRCA* testing inside the onco-genetic framework marks an important step-forward in BC management. However, the integration of somatic testing, digital pathology, and artificial intelligence-driven models could refine patient selection for tailored treatments.

Introduction

Germline pathogenic variants (gPV) in *BRCA1* and/or *BRCA2* genes are associated with an increased risk of developing breast cancer and ovarian cancer [1]. *BRCA2* gPV may also increase the risk of male breast cancer with an estimated lifetime risk of approximately 13 % for male carriers. Other genes associated with increased breast cancer risk include *TP53*, *PTEN*, *CDH1*, *STK11*, *MSH6*, *BARD1*, *ATM*, *CHEK2*, *RAD51C*, *RAD51D* and *PALB2* [2,3]. With the introduction of poly (ADP-ribose) polymerase (PARP) inhibitors firstly in advanced and then in early-stage settings, a new approach to genetic counselling – the Mainstreaming Cancer Genetics (MCG) – has been introduced into the clinical practice. This model enables medical oncologists and others specialists involved in breast cancer diagnosis and treatment to directly refer patients for *BRCA* genetic testing when it has therapeutic implications. While somatic testing is increasingly being explored as a potential screening tool for *BRCA* gPV detection, current supporting evidence remains limited (level IIB in the ESCAT score) [4]. Therefore, peripheral blood analysis continues to be the standard reference test.

This document aims to provide an update on predictive *BRCA* testing and support its implementation in clinical practice, with a focus on the following patient groups:

1. Patients with clinical characteristics associated with an increased likelihood of *BRCA1/2* gPV, regardless of family history.
2. Patients without clinical characteristics associated with an increased likelihood of *BRCA1/2* gPV but who are eligible for specific treatments if a gPV is detected.

3. Patients with family history suggestive of a high likelihood of *BRCA1/2* gPV.

Furthermore, the document also addresses surgical and radio-therapeutic treatments in patients already affected by breast cancer.

Methods

Document development and composition of the multidisciplinary working group

The document has been developed, discussed, reviewed, and approved by a multidisciplinary Expert Panel composed of 33 clinicians representing major Italian scientific associations of Italian Association of Medical Oncology (AIOM), Italian Association of Radiotherapy and Clinical Oncology (AIRO), Italian National Association of Breast Surgeons (ANISC), Italian Society of Pathological Anatomy and Diagnostic Cytology (SIAPeC-IAP), Italian Society of Surgical Oncology (SICO), Italian Society of Human Genetic (SIGU), Italian Society of General Practice (SIMG), Italian Society of Medical and Interventional Radiology (SIRM). It is based on expert consensus and incorporates the best available evidence, aligning and harmonizing the current eligibility criteria for *BRCA* testing in accordance with Italian National Guidelines and Clinical Recommendations.

The working group encompassed medical, surgical, and radiation/clinical oncologists; medical and molecular geneticists; clinical molecular biologists; surgical and molecular pathologists; organ specialists such as gynecologists, gastroenterologists and urologists, and pharmacologists; methodologists.

Grading of Recommendations, Assessment, Development and Evaluation (GRADE) recommendations

After identifying specific outcomes of benefit and harm deemed critical for decision-making, the GRADE approach was applied to assess the certainty of evidence for each selected outcome, evaluating five key domains: study limitations, imprecision, indirectness, inconsistency, and publication bias.

The overall quality of evidence, which forms the basis for the final recommendations, is classified into four levels: high, moderate, low, or very low.

The strength of the final recommendation is determined through a voting process, with four possible outcomes: strong in favor, conditional in favor, conditional against, and strong against.

The clinical recommendation reflects the practical clinical significance of the procedure.

In the [Appendix A](#) are presented literature search, PRISMA flow diagram and evidence to decision framework for the GRADE query on contralateral prophylactic mastectomy.

Results

Eligibility criteria for onco-genetic counselling

The criteria for onco-genetic counselling in breast cancer patients are outlined below and in [Table 1](#). The pathway for genetic testing in breast cancer patients should be tailored according to the “a priori” likelihood of gPV and the primary objective of testing, whether predictive (for treatment decision) or preventive” (for risk assessment and management).

Patients with clinical characteristics associated to an increased likelihood of harboring a BRCA1/2 gPV, regardless of family history

- woman with breast and ovarian cancer;
- woman with breast cancer diagnosed at ≤ 40 years;
- woman with triple negative breast cancer (TNBC) diagnosed at any age;
- woman with bilateral breast cancer diagnosed at ≤ 50 years;

Table 1

Eligibility criteria for performing germline BRCA genetic testing.

Patients with clinical characteristics associated to an increased likelihood of BRCA1/2 gPV, regardless of family history

Woman with breast and ovarian cancer
 Woman with breast cancer diagnosed at ≤ 40 years
 Woman with triple negative breast cancer (TNBC) diagnosed at any age
 Woman with bilateral breast cancer diagnosed at ≤ 50 years
 Man with breast cancer

Patients without clinical characteristics associated to an increased the likelihood of BRCA1/2 gPV, but eligible for specific treatments in case a gPV is identified:

Early breast cancer patient with hormone receptor-positive (HR+), HER2-negative (HER2-) disease and ≥ 4 lymph nodes involved
 Breast cancer patient with HR+/HER2- disease, with residual disease after neoadjuvant chemotherapy and a CPS/EG score (clinical and pathologic stage/estrogen receptor status and histologic grade) ≥ 3 ;
 Metastatic breast cancer patient with HR+/HER2- disease, previously treated with anthracycline and/or taxane-based chemotherapy (or deemed ineligible for it), progressing after Cyclin D Kinase 4/6 inhibitors (CDK4/6i) for advanced disease

Patients with family history associated with a high likelihood of BRCA1/2 gPV

BC patient diagnosed between 41 and 50 years with a first degree relative* with:

- Breast cancer diagnosed before the age of 50 years
- Ovarian cancer (excluding mucinous and borderline histologies), diagnosed at any age
- Bilateral breast cancer
- Pancreatic cancer
- Prostate cancer

Breast cancer patient diagnosed after the age of 50 years with two first-degree relatives* (related to each other), including one who is a first-degree relative of the patient, affected by breast, ovarian, or pancreatic cancer.

Presence of a gPV already identified in a relative.

- man with breast cancer.

For this group of patients, BRCA testing can be requested as a fast-track procedure to guide the most appropriate treatment decisions, both locoregional and systemic, based on the test results. This request can be made not only by geneticists but also by oncologists and surgeons, at a multidisciplinary team (MDT) level discussion, provided they have received adequate training to ensure that patients receive accurate and comprehensive information. Whenever a gPV or a variant of unknown significance (VUS) is identified, genetic counselling must be promptly activated. Conversely, in patients who do not carry a gPV or VUS, testing for other predisposing genes may be considered (either as part of the same multigene panel or at a later stage).

AIOM guideline update has revised the diagnostic criteria for all breast cancer cases, raising the age threshold from 36 to 40 years at the time of diagnosis. This adjustment aligns with the increased prevalence of gPV observed in this age group [5–7]. To assess the feasibility of this amendment, the Italian Cancer Registry estimated the number of breast cancer diagnoses in Italy among patients aged 36–40 years, identifying approximately 1,800 additional patients eligible for testing [8]. Based on the current capacity of national laboratories, this increase in BRCA testing volume was deemed sustainable. Additionally, the age criterion—previously set at 60 years—has been removed for patients diagnosed with TNBC. As a result, TNBC itself is now considered a standalone indication for BRCA testing, regardless of the patient’s age at diagnosis [9].

Patients without clinical characteristics suggesting a high likelihood for BRCA1/2 gPV but eligible for specific treatments if a gPV is detected:

- Any patient affected by hormone receptor-positive (HR+), HER2-negative (HER2-) early breast cancer and ≥ 4 lymph nodes involved;
- Any patient affected by HR+/HER2- early breast cancer, with residual disease after neoadjuvant chemotherapy and a clinical and pathological stage/estrogen receptor status and histologic grade (CPS/EG) score ≥ 3 ;
- Any patient affected by advanced/metastatic HR+/HER2- breast cancer, previously treated with anthracycline and/or taxane-based chemotherapy (or deemed ineligible for it), progressing after

* First-degree relative (parent, brother/sister, son/daughter) with breast cancer, ovarian cancer or pancreatic cancer. In the paternal lineage, second-degree relatives (grand-mother, aunts) must be considered.

cyclin-dependent kinase 4 and 6 inhibitors (CDK4/6i) for advanced disease.

For this patient category, *BRCA* testing is requested for therapeutic purposes (to assess eligibility for PARP-inhibitors) by a trained oncologist, ensuring that patients receive accurate and comprehensive information. Genetic counselling must be promptly activated if a gPV or VUS is detected.

Patients with family history associated with a high likelihood of BRCA1/2 gPV

- a) Breast cancer patient diagnosed between 41 and 50 years with a first degree relative affected by:
 - Breast cancer, diagnosed within the age of 50 years;
 - Ovarian cancer, excluding mucinous and borderline histologies, diagnosed at any age;
 - Bilateral breast cancer;
 - Pancreatic cancer;
 - Prostate cancer.
- b) Breast cancer patient diagnosed after the age of 50 years with two first degree relatives (related to each other, including one who is first-degree relative of the patient), affected by breast, ovarian or pancreatic cancer.
- c) Presence of a gPV already identified in a relative.

For patients who may have a paternally inherited *BRCA* gPV, due to the male interposition, the second-degree relatives can be considered as first-degree ones.

For this patient category, traditional genetic counselling is activated to optimize the genetic evaluation. This process involves verifying tumor characteristics in affected relatives, assessing their distribution within the family tree, determining the appropriateness of genetic testing, and identifying the most informative index case. A comprehensive evaluation should also include differential diagnosis among other hereditary syndromes, cancer risk estimation, and prevention strategies in case of uninformative results. In addition, co-segregation and *in silico* analyses should be considered to better interpret VUS

When *BRCA* genetic testing is required for therapeutic decisions, genetic counselling should be conducted within one week of the request. If geneticists are unavailable, a trained medical or clinical/radiation oncologist or surgeon may request the test ensuring that patients receive accurate information. The results must then be reviewed by the MDT, including a geneticist, to ensure proper interpretation.

The *BRCA* test must be integrated into a structured genetic pathway, conducted by health providers with both oncological and genetic expertise. These skills are essential for both pre-test and post-test counselling. A predictive *BRCA* testing for therapeutic decisions can be prescribed not only by geneticist, but also by oncologists and breast surgeons with oncological skills, who are responsible for providing adequate information on the implications of the results.

The test results can be:

- Informative: the predisposing variant has been identified;
- Uninformative: the predisposing variant has not been detected, but its presence cannot be entirely excluded, or a VUS has been identified with no confirmed clinical significance.

Germline pathogenic variants are categorized into five classes according to the Evidence-based Network for the Interpretation of Germline Mutant Alleles (ENIGMA) classification:

- Category 1: benign = no clinical significance
- Category 2: likely benign = unlikely to be associated with disease.
- Category 3: variant of unknown significance = Insufficient evidence to determine clinical relevance.

- Category 4: likely pathogenic = strong evidence suggests an association with disease.
- Category 5: pathogenic = confirmed to be disease-causing

A true negative result occurs only when a pathogenic variant previously identified in a family member is not detected in the tested subject. The test can be extended to other family members, who wish to undergo testing, only in case of an informative result, starting at 18 years of age. *BRCA* test in minors is currently not recommended, as the associated risk begins to increase in adulthood. In case of VUS, extending the test to other family members is not indicated outside of research projects. Surveillance programs should be based on family history and other established risk factors.

Positioning of predictive gBRCA1/2 test for the use of PARP inhibitors

The availability of PARP inhibitors in both early and advanced settings for patients with HER2-negative breast cancer carrying a *BRCA1/2* gPV [10–12] has necessitated a revision of the *BRCA* testing framework, as shown in Fig. 1.

The presence of *BRCA* gPV has crucial implications at multiple levels. From a surgical perspective, *BRCA* gPV patients should be offered bilateral mastectomy (even in case of radiological complete response after neoadjuvant treatment). From a medical standpoint, 1 year of adjuvant olaparib represents a priority treatment option for patients at high-risk of relapse, as shown in Fig. 1.

In patients with early TNBC, *BRCA* testing should be performed before initiating any treatment, whether in the neoadjuvant or adjuvant setting. Within 7 days, the MCG should be conducted by an oncologist or surgeon, along with blood sample collection for the *BRCA* test. A psychologist can be involved upon request. The genetic analysis should be completed within a maximum of 3 weeks and the final result should be released within 2 days. In case of a positive result, the discussion should take place in the presence of a breast surgeon and an oncologist, along with a plastic surgeon and psychologist, as needed.

In TNBC with a radiological diameter of less than 1 cm, surgery should be the first therapeutic choice and bilateral mastectomy should be considered. If after primary surgery the pathological stage is $\geq$ II, adjuvant chemotherapy should be administered for at least six cycles followed by Olaparib therapy for one year. In case of bilateral prophylactic mastectomy an early plastic surgery consultation is warranted to plan in advance the best reconstructive options to be shared with the patient.

In tumours with a clinical diameter between 1 and 2 cm, neoadjuvant treatment with anthracyclines, cyclophosphamide and taxane (with or without platinum-derived drugs) may be considered. In stage II-III tumours at diagnosis, pembrolizumab combined with chemotherapy (carboplatin and paclitaxel for 4 cycles followed by anthracycline and cyclophosphamide for 4 cycles) should be the standard of care according to the results of the Keynote-522 study [13]. However, it should be discussed with the patient the opportunity to receive only neoadjuvant chemotherapy without pembrolizumab, followed by Olaparib in case of residual disease, due to a limited benefit in pathological complete response (pCR) rate from the addition of pembrolizumab (69.1 % vs. 62.1 % in the placebo arm). Furthermore, the Olaparib treatment in adjuvant setting shows a better toxicity profile compared to pembrolizumab and achieves an OS advantage in 4 years follow-up period compared to 75 months in the Keynote-522 study.

For patients who are undergoing neoadjuvant chemotherapy and carry a *BRCA* gPV, bilateral mastectomy with immediate reconstruction should be discussed and considered, even in cases of radiological complete response. If residual disease is present, adjuvant Olaparib should be prioritized over pembrolizumab (see Fig. 2).

In patients with HR+/HER2- early breast cancer who do not present clinical characteristics suggesting a high likelihood of *BRCA1/2* gPV, predictive *BRCA* testing should be offered to those eligible for one year




Fig. 1. Possible treatment pathways for gBRCA test in early and advanced breast cancer, either Triple Negative (A) and HR+/HER2- (B) BRCA+: BRCA positive; NEO/ADJ: Neoadjuvant/Adjuvant; CT: Chemotherapy; ICI: Immune Checkpoint Inhibitors; PD-L1+: PD-ligand 1 positive; PARPi: Poly (ADP-ribose) polymerase inhibitors; PD-L1-: PD-ligand 1 negative; BC: breast cancer; HR+:Hormone Receptors positive; HER2-: human epidermal growth factor receptor 2 negative; ET: endocrine therapy; MBC: Metastatic Breast Cancer; CDK4/6i: Cycline D Kinase 4/6 inhibitors; RT: Radiotherapy.




Fig. 2. Multidisciplinary pathway of Triple Negative Breast Cancer at stage I-III diagnosis MCG: Mainstreaming Cancer Genetics; BRCA+: BRCA positive; BPM: Bilateral Prophylactic Mastectomy; ADJ: Adjuvant; CT: Chemotherapy; PARP-i: Poly (ADP-ribose) polymerase inhibitors; NEOADJ: Neoadjuvant; ICI: Immune Checkpoint Inhibitors; pCR: pathological Complete Response, RT: Radiotherapy.

of adjuvant olaparib (in combination with adjuvant endocrine treatment), based on the OlympiA trial eligibility criteria. In these cases, the BRCA test should be performed within 4 weeks of surgery or during adjuvant chemotherapy to ensure results are available within the required timeframe to start olaparib within 12 weeks of completing local treatment (see Fig. 3).

In the metastatic setting, predictive BRCA testing should be performed at the time of diagnosis of HER2- advanced disease given the availability of olaparib and talazoparib as treatment options. More specifically, in patients with advanced TNBC, PARP inhibitors (olaparib or talazoparib) represent a valuable first-line treatment option in case of PD-L1 negative status. In patients with PD-L1 positive tumors, who receive chemotherapy plus immunotherapy in the first line setting, PARP-inhibitors may be considered from the second-line onward. According to the Italian label, both olaparib and talazoparib should be preceded by treatments with platinum-derived drugs, anthracycline and/or taxanes in any disease setting, unless these are contraindicated

(Fig. 1A).

In advanced HR+/HER2- breast cancer, the only PARP-inhibitor currently approved by the Italian drug agency (AIFA) and reimbursed by the national health system is talazoparib. This agent represents a key treatment option in endocrine-resistant patients with BRCA gPV who have already received endocrine-based therapy, including CDK4/6 inhibitors. Additionally, patients must have previously undergone anthracycline-taxane based chemotherapy either in the early or advanced setting (Fig. 1B).

Treatment of patients with BRCA1/2 gPV and breast cancer: focus on the role of bilateral mastectomy/contralateral prophylactic mastectomy/bilateral salpingo-oophorectomy

The optimal surgical approach – breast-conserving surgery (plus radiation therapy [RT]) vs radical mastectomy in BRCA1/2 carriers affected by breast cancer is still debated. Several meta-analyses have




Fig. 3. Multidisciplinary pathway of HR+/HER2- breast cancer at stage I-III diagnosis HR+: Hormone Receptors positive; HER2-: human epidermal growth factor receptor 2 negative; NEO/ADJ: Neoadjuvant/Adjuvant; CT: Chemotherapy; ET: endocrine therapy; MCG: Mainstreaming Cancer Genetic; BRCA+: BRCA positive; PARP-i: Poly (ADP-ribose) polymerase inhibitors; BC: Breast Cancer; pCR: pathological Complete Response; CPS-EG: clinical and pathologic stage and estrogen receptor status and histologic grade; RT: Radiotherapy.

reported that the risk of a second ipsilateral breast cancer is approximately 24 % after 15 years in patients treated with conservative surgery plus RT compared to those undergoing mastectomy [14,15]. However, this increased risk did not impact the mortality rate at 15 years [16]. The use of prophylactic contralateral irradiation has been evaluated in a phase II case/control trial showing a 80 % reduction in breast cancer risk [17].

Compared to non-BRCA carriers, the risk of contralateral breast cancer 25 years after the initial diagnosis is increased by 53 % for BRCA1 and 65 % for BRCA2 gPV carriers. Furthermore, the cumulative ovarian cancer risk to age 80 years is 44 % for BRCA1 and 17 % for BRCA2 carriers [18]. The contralateral breast cancer risk remains elevated also in patients who have undergone bilateral prophylactic salpingo-oophorectomy and is influenced by the age at first diagnosis [19]. Given the known increased risk of contralateral breast cancer, many women choose to undergo bilateral mastectomy at the time of breast cancer surgery rather than in a later stage [20]. This underscores the importance of performing the BRCA test at the time of breast cancer diagnosis (when criteria for BRCA testing are met), ensuring that results are available within 4 weeks. Some retrospective evidence suggests that contralateral prophylactic mastectomy (CPM) in BRCA1/2 gPV carriers may reduce mortality, mostly in the second decade after surgery [21–23].

Particular attention should be given to young patients. Women aged

≤40 years who have already been diagnosed with breast cancer and carry a BRCA gPV are at higher risk of developing a second malignancy over their lifetime. In the group of women with early-stage breast cancer, metachronous ovarian cancer is a common cause of mortality [24]. In these patients, to undergo bilateral risk-reducing mastectomy and/or salpingo-oophorectomy appears to have a positive impact on prognosis, leading to lower rates of recurrence, secondary breast and/or ovarian cancers, and mortality compared to those who do not pursue these strategies. Recent data from a cohort study of BRCA1/2 gPV, show that oophorectomy among patients already affected by breast cancer reduces the specific mortality risk of 56 % (HR = 0.44; 95 % CI, 0.25–0.78; P = 0.005), mostly in BRCA2 patients (HR = 0.22; 95 % CI, 0.06–0.84; P = 0.03) [25].

To provide recommendations based on the GRADE system, the following clinical question was voted on by the panelists, as reported in Table 2:

For this GRADE query the research evidence is the following:

Breast cancer is the leading cause of cancer-related death among women worldwide, being 522,000 the patients deceased in 2012 (15 % of all female deaths) [26]. Furthermore, it represents the most common cancer in all countries with 1,7 million of women affected [27].

In Italy about 58,000 new breast cancer/year are estimated, among those 5 % arise in genetically predisposed women with BRCA1/2 gPV. Since patients with BRCA1/2 gPV are at increased risk of ovarian and

Table 2

GRADE clinical question on CPM versus no CPM in patients with a history of surgically treated breast cancer and a BRCA gPV.

Is contralateral prophylactic mastectomy indicated in patients with a history of surgically treated breast cancer and BRCA gPV?

RECOMMENDATION: In patients with operated breast cancer and gPV BRCA, contralateral prophylactic mastectomy may be performed.

Strength of the recommendation: CONDITIONAL IN FAVOUR

Reasons/Comments to the benefit/risk ratio:

Outcome of benefit. The panel deemed the following outcomes of benefit as critical:

Breast cancer mortality, disease free interval (DFI), breast cancer incidence, quality of life, Patient Reported Outcomes (PROs). Supporting evidence comes from observational studies indicating that contralateral prophylactic mastectomy was associated with a significant improvement of: breast cancer mortality (HR 0.46; IC95% 0.20–0.71), breast cancer incidence (OR 0.08; IC95% 0.03–0.19) DFI (HR 0.42; IC95% 0.26–0.69) compared to no procedure. The procedure was well tolerated with minimal impact on quality of life, as indicated by PROs. Considering this evidence, the impact of the procedure on these benefit outcomes was rated as great by 9 and moderate by 2 panelists (totally 11 voters). The assessment was affected by the low quality of evidence, primarily due to the following biases: study design limitations, including selection, detection, recall, and attrition biases, as well as result imprecision.

Outcome of harms. The panel identified the following outcomes of harms: long terms coping with cosmetic results, early and late physical morbidity. Five observational studies reported on the effects of contralateral prophylactic mastectomy effects on long term cosmetic results. None of the studies reported on early and late physical morbidity. Due to the low quality of evidence, the impact of the procedure on the outcomes of harms was considered moderate by all voters (totally of 11 voters)

Given the limitations of the included observational studies and the low quality of evidence, the panel assessed the balance between benefits and harms as uncertain in favor of the contralateral prophylactic mastectomy.

Future research implications: perspective studies are needed.

Proofs quality: Outcome of benefit: Very low; Outcome of harms: Very low

Global proofs quality: VERY LOW

Declaration of competing interest (COI): no COI to declare

breast cancer, contralateral too, CPM could be considered to prevent the second breast cancer onset. As preventive measure, risk reducing mastectomy remains controversial. Potential benefits include reduction of breast cancer risk and increased psychological wellbeing. Potential disadvantages regard procedure invasiveness and subsequent morbidity, such as reduction of body image satisfaction and reduction of breast tactile sensitivity. It remains clear that this surgery must be performed in high-volume centres with specific experience.

Discussion

BRCA testing is a key component in the management of early and metastatic HER2-negative BC, as the presence of *BRCA* gPV currently serves as a theragnostic biomarker for access to PARP inhibitors. In Italy, the national health system provides *BRCA* testing free of charge, accordingly to National Guidelines. Thus, defining eligibility criteria for *BRCA* test is essential to ensure treatment approval and reimbursement. Beyond the new implications arising from the recent introduction of PARP inhibitors, the predictive role of genetic testing remains well-established in guiding surgical decision-making. In fact, offering bilateral mastectomy at the surgical time seems to provide an overall survival (OS) benefit compared to unilateral mastectomy [28]. Furthermore, the rate of bilateral mastectomy in breast cancer patients is around 75 % when the *BRCA* result is known before the surgery, compared to 15 % when is released after [29,30].

The introduction of MCG, performed by health care providers such as surgeons or oncologists, ensures that test results are delivered within an appropriate timeframe, allowing for optimal medical treatment planning.

On the medical treatment point of view, knowing the *BRCA* status, allows to offer one-year treatment with Olaparib, that provides OS benefit compared to placebo arm [31]. We well know that TNBC patients treated with neoadjuvant therapy according to the Keynote-522 trial derive a benefit in OS by the adjuvant pembrolizumab, mostly in case of Residual Cancer Burden (RCB)-2 [32,33] but not in case of RCB-1 or RCB-3. In Italy, where the Pembrolizumab plus Olaparib combination is not permitted, knowing the *BRCA* status can aid to choose the right adjuvant strategy.

In HR+ tumors, the alternative to Olaparib could be Abemaciclib or Ribociclib, that did not show OS benefit compared to endocrine treatment alone yet. Some retrospective data in the metastatic setting seem to suggest a worse prognosis of gPV *BRCA2* patients treated with CDK4/6i compared to wild-type *BRCA* [34]. However, given the small number of gPV *BRCA* patients included in these studies, caution is warranted in drawing definitive conclusions about potential resistance mechanisms induced by drug exposure and the optimal treatment sequence. Further studies are needed to address these important questions.

By this way, our guidelines assure the *BRCA* test to all patients responding to eligibility criteria for early and metastatic breast cancer, regardless the family history or the patient characteristics'.

In the case of indications for RT (both after breast-conserving surgery and mastectomy) and adjuvant PARP-i or CDK4/6i, there is no recommendation to administer concomitant RT outside the context of a clinical trial, due to the potential increase in toxicity and the lack of robust data on combination treatments. The consensus statement endorsed by the European Society for Radiotherapy and Oncology (ESTRO) recommends starting adjuvant PARP-i or CDK4/6i after the completion of the postoperative RT course [35].

The MCG process has enabled a more efficient redistribution of resources, optimizing the pathway at a time when the demand for genetic testing is steadily increasing, while ensuring the appropriate level of oversight and support for patients with a positive result.

The new *BRCA* test paradigm necessarily needs to be integrated in a multidisciplinary medical practice. Such an approach would not only promote adherence to test criteria, but also facilitate the monitoring of target treatments. There is widespread consensus on the crucial role of

the MDT in discussing the timing of the test, taking into account the unique characteristics of each patient and their disease. To allow proper effective discussion at MDT meetings especially in the pre-operative scenario all experts of the MDT should be fully aware of the MCG process and promptly request the test when seeing patients at the beginning of their diagnostic analyses. This would allow to mitigate the turn around time of *BRCA* testing and to obtain the information in a timely way. Finally, the role of psychosocial support is part of the *BRCA* test since depression, anxiety, stress, social and economic challenges, should be assessed and addressed for a comprehensive counselling, mostly in case of young patients who are worried for their kids.

In this context, the AIOM Guidelines Panel identified the redefinition of *BRCA* testing eligibility criteria as a fundamental and urgent priority. One of the most debated topics within this working group was the potential incorporation of the universal genetic testing into the Italian Breast Cancer Guidelines, an approach that has been endorsed since 2019 by the Consensus Guidelines on Genetic Testing for Hereditary Breast Cancer from the American Society of Breast Surgeons [36]. While universal genetic testing may be considered an ideal strategy, its implementation faces several critical challenges, including: costs and, in general, sustainability within publicly funded healthcare systems, scarcity of genetic counselling services, increased detection of VUS, limited utility of *BRCA* testing in patients diagnosed with breast cancer at ≥ 65 years of age, especially for HR+/HER2- subtype. Among these challenges, sustainability is particularly crucial in Italy, where a universal healthcare system ensures equitable access to medical services, but demands strategic resource allocation. A cost-effectiveness analysis demonstrated that universal *BRCA* testing for breast cancer patients becomes cost-ineffective when the test cost exceeds a certain threshold [37]. To improve cost-effectiveness, it is essential to increase the number of tests processed per laboratory, centralizing genetic testing in high-volume Hub centers while ensuring a regionally balanced distribution of services. In the Emilia-Romagna region, where 4500 new breast cancer at year are estimated, 4 Hub centers are able to provide all predictive *BRCA* test, accordingly to National guidelines, corresponding to about 53 % of all new breast cancer diagnosed [38]. As reported by Cortesi et al., an intensive surveillance program has been provided, with annual mammography, combined with semi-annually ultrasound $\pm$ annual magnetic resonance imaging, accordingly to moderate and high risk for breast cancer development, calculated by the Tyrer-Cuzick model [39]. By this, we can ensure an adequate follow-up for at risk people, even in the absence of a *BRCA* gPV.

While universal *BRCA* testing is likely the future direction, at present, this approach is not considered sustainable within the Italian healthcare system. Therefore, the AIOM multidisciplinary guidelines panel opted for a gradual expansion of *BRCA* testing eligibility rather than an immediate shift to universal testing—an approach that will eventually become inevitable.

This progressive transition has been facilitated through three key modifications: abolition of the age threshold for TNBC, making it a standalone criterion for *BRCA* testing, raising the age threshold for non-TNBC cases from 36 to 40 years and formal introduction of predictive *BRCA* testing in patients who, despite lacking a strong personal or family history, suggestive of a high mutation probability, may still be eligible for specific targeted therapies, particularly PARP inhibitors.

The low percentage of TNBC among the breast cancer subtypes (15–20 %) made possible to test all patients belonging to this category, having the highest gPV *BRCA* rate (13–15 %). Otherwise, testing all HR+/HER2- breast cancer, that represent the vast majority of cases, would have meant unsustainable costs for our Public Health System, given the low mutation rate in this population (5–6 %). Similarly, the low rate of gPV *BRCA* in HER2+ subtype does not justify extending testing to all those patients. Nowadays, we think that this incremental expansion of *BRCA* testing criteria ensures greater accessibility to precision oncology treatments while maintaining sustainability within the current healthcare framework. As testing costs become more affordable,

we intend to extend the analysis to all breast cancer patients. This will also allow us to expand the test indication to patients with multiple tumors in addition to breast cancer, especially in case of association with pancreatic carcinomas.

The introduction of MCG for predictive *BRCA* testing has proven effective in reducing the reliance on geneticists, limiting specialist genetic services primarily to patients with significant and suggestive family/personal history. A prospective study demonstrated that MCG provided by surgeons or nurses delivered sufficient information for most breast cancer patients to make informed decisions about genetic testing, with minimal distress, compared to traditional geneticist-led pre-test counselling [40]. The only statistically significant difference regarded the less frequent discussion about second breast cancer or ovarian cancer in the MCG compared to the traditional genetic counselling. However, in both groups, genetic knowledge and patient satisfaction levels were comparable.

Nevertheless, genetic counselling remains essential in cases of gPV or VUS, for cascade testing in relatives, for a comprehensive discussion of surgical options and ovarian cancer risk, and for further risk assessment in other inherited cancer syndromes in patients with uninformative results. Unlike predictive testing, post-test genetic counselling in these cases can be conducted without strict time constraints. Nonetheless, a recent survey conducted across the Canadian healthcare system revealed that only 67 % of patients informed their relatives about a positive *BRCA* test, highlighting the need for additional support in result dissemination [41].

Albeit our guidelines have been focused on *BRCA1/2* testing, they do not preclude the test for other hereditary syndromes genes, (i.e. *TP53*, *PTEN*, *STK11* and *CDH1*), that would impact management decisions such as CPM, or avoiding the use of radiation. At this aim, patients undergone predictive analysis without carrying *BRCA1/2* gPV, can receive testing for other predisposing genes, as part of the same multigene panel or at a later stage, accordingly to specific personal or family history.

The increasing use of multigene panel testing allows for the identification of additional high- and moderate-penetrance genes associated with hereditary breast cancer at the time of initial breast cancer diagnosis. It has been estimated that 8–15 % of patients testing negative for *BRCA1/2* carry a pathogenic variant in other hereditary cancer genes [42,43]. This expanded genetic profiling not only accelerates the diagnosis of other hereditary syndromes but also facilitates cascade testing in affected families. Furthermore, specific genetic findings can have direct clinical implications, as in the case of *PALB2* gPV, where surgical management aligns with that used for *BRCA1/2* gPV carriers and is also expected to be incorporated into the access criteria for PARP-inhibitors [44]. However, the widespread adoption of multigene panel testing also increases the detection of VUS, which can pose significant challenges. Although, at the beginning of multigene panel adoption, patients could misinterpret uncertain results, developing inaccurate cancer risk perceptions, experiencing prolonged distress despite psychosocial support, or even pursuing prophylactic surgeries inconsistent with current guidelines [45–47], the increasing knowledge of VUS meaning and the improvement into the right choice of genes to include in a test, provided better information on correct cancer-related surgical and screening management. Indeed, more recent data show that VUS carriers received less CPM, prophylactic mastectomy and oophorectomy and less annual breast magnetic imaging resonance compared to gPV ones [48–50]. From a healthcare perspective, the increased detection of VUS also places a burden on genetic services, due to long-term variant reclassification, and recontact protocols. Effective VUS communication involves clarifying the actual probability of future pathogenic reclassification [51]. However, the timeline for VUS reclassification remains undefined, presenting an ongoing challenge for both clinical laboratories and healthcare services. A recent cohort study analyzing 94,453 variants across 3,272,035 individuals reported that 4.5 % of VUS in hereditary disease genes were reclassified over an 8-year period, emphasizing the

extensive workload and the role of emerging machine learning-based computational methods in variant resolution [52].

Concerns remain regarding the utility of genetic testing in patients diagnosed with HR+/HER2– breast cancer without family history at ≥65 years, given the high negative predictive value (>99 %) and low positive rate (1.09 %) in this population [53,54].

Emerging technologies, including somatic *BRCA* testing and artificial intelligence (AI)-based digital pathology, could serve as pre-screening tools, requiring confirmation through direct sequencing in peripheral blood. Some studies suggest that AI can accurately predict homologous recombination deficiency, with Relative Operating Characteristic (ROC) curves of 0.86 and 0.81, particularly in HR+/HER2– Breast cancer subtypes [55,56].

Interestingly, the European Society of Medical Oncology (ESMO) Precision Medicine Working Group supports the use of tissue Next Generation Sequencing (NGS) as a substitute for germline *BRCA1/2* testing in patients with endocrine-resistant HR+/HER2– metastatic breast cancer, given the high accuracy of tissue sequencing. However, reflex germline testing is still recommended in cases where no tumor alterations are detected but the probability of harboring a germline *BRCA1/2* gPV remains high, considering that NGS may fail to detect approximately 7 % of germline *BRCA* gPV[4].

Despite these recommendations, regulatory barriers still limit the full adoption of this approach in Italy, where access to PARP inhibitors is restricted to patients with a *BRCA* gPV identified through germline testing, in accordance with the registration trials. Consequently, implementing NGS as a pre-screening tool for drug eligibility would require a more flexible interpretation of current regulatory criteria, a topic that warrants further discussion at the decision-making level in future policy frameworks.

To conclude, the integration of predictive *BRCA* testing into the onco-genetic framework represents an important step-forward, driven by the constantly evolving therapeutic landscape of breast cancer. This process has required epidemiological assessments, structural optimizations and cost-effectiveness evaluations to ensure feasibility within the healthcare system. Despite these advancements, several open questions remain. Notably, even with the incremental expansion of *BRCA* testing criteria, a small subset of patients may still be overlooked, potentially missing the opportunity to access effective treatments such as PARP inhibitors. Additionally, while these targeted agents have been transformative for HER2– *BRCA*-related breast cancer, several alternative treatment options are available in the same clinical settings (i.e., pembrolizumab/capecitabine for early TNBC or abemaciclib for HR+/HER2– early breast cancer), thus necessitating careful considerations at a single case level. Finally, as precision oncology continues to evolve, the integration of somatic testing, digital pathology, and artificial intelligence-driven models may further refine patient selection, optimizing treatment allocation and ensuring that innovative therapies reach those most likely to benefit.

Funding sources

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Cortesi Laura reports a relationship with AstraZeneca Pharmaceuticals LP, Merck Sharp & Dohme Corp, Daiichi Sankyo Inc and Pfizer Inc that includes: board membership and consulting or advisory. Cortesi, Laura reports a relationship with Novartis, Gilead Sciences Inc, Roche Diagnostics Corporation that includes: speaking and lecture fees. Miglietta Federica, reports a relationship with Roche, Novartis, Pfizer/Seagen, MSD, Astrazeneca, Daiichi Sankyo, Menarini Stemline, Gilead that

includes: speaking and lecture fees. Biganzoli Laura, reports a relationship with Amgen, AstraZeneca, Boehringer-Ingelheim, Daiichi-Sankyo, Eisai, Exact Sciences, Gilead, Lilly, Menarini, MSD, Novartis, Pfizer, Pierre Fabre, Roche, Sanofi, SeaGen that includes: board membership and consulting or advisory. Biganzoli Laura, reports a relationship with Celgene, Genomic Health, Novartis that includes: Institutional financial interests. Biganzoli Laura, reports a relationship with AstraZeneca, Daiichi-Sankyo, MSD that includes: Travel reimbursement. Del Mastro Lucia reports a relationship with Eli Lilly; Novartis; Roche; Daiichi Sankyo; Seagen; Astrazeneca; Gilead; Pierre Fabre; Pfizer.

Menarini Stemline; MSD that includes: board membership and consulting or advisory. Del Mastro Lucia reports a relationship with Roche, Novartis, Pfizer, Eli Lilly, Astrazeneca, MSD, Seagen, Gilead, Pierre Fabre, Eisai, Exact sciences, Ipsen, GSK, Agendia, Menarini Stemline that includes: speaking and lecture fees. Del Mastro Lucia reports a relationship with Roche, Daiichi Sankyo, Astrazeneca, Gilead, Menarini Stemline that includes: Travel reimbursement. Foglietta Jennifer reports relationship with Menarini Stem Line that includes: board membership and consulting or advisory. Foglietta Jennifer reports relationship with Novartis that includes: speaking and lecture fees. Foglietta Jennifer reports relationship with Roche, Sophos, Pfizer that includes: travel reimbursement. Marchio' Caterina reports a relationship with Roche, Astrazeneca, Daiichi Sankyo, Menarini, Illumina that includes board membership and consulting or advisory. Musolino Antonino reports a relationship with Eli Lilly, Eisai Europe, Seagen, Daiichi-Sankyo, AstraZeneca, Novartis that includes board membership and consulting or advisory. Musolino Antonino reports a relationship with Eli Lilly and Roche that includes: funding or grants. Musolino Antonino reports a relationship with Pfizer that includes: travel reimbursement. Zambelli Alberto reports a relationship with Pfizer, Lilly, Novartis, Roche, AstraZeneca, DaiichiSankyo, Seagen, ExactSciences, MSD, Gentili that includes: board membership and consulting or advisory. Zambelli Alberto reports a relationship with Roche, DaiichiSankyo, AstraZeneca, Novartis that includes: travel reimbursement. Curigliano Giuseppe reports a relationship with advisory board from Roche, Novartis, Lilly, Pfizer, Astra Zeneca, Daichii Sankyo, Ellipsis, Veracyte, Exact Science, Celcuity, Merck, BMS, Gilead, Sanofi, Menarini that includes: board membership and consulting or advisory. Dessena Massimo reports a relationship with Novartis that includes: board membership and consulting or advisory board. Fabi Alessandra reports a relationship with Roche, Lilly, Novartis, AstraZeneca, Pfizer, Seagen, Gilead, MSD, Menarini Stemline, Dompè Biotech that includes: board membership and consulting or advisory. Fabi Alessandra reports a relationship with Astra Zeneca, Roche, Lilly, Novartis, Gilead, Pfizer, Daiichi Sankyo Exact Sciences that includes: speaking and lecture fees. Fabi Alessandra reports a relationship with Astra Zeneca, Roche that includes: funding or grants to the Fondazione Policlinico Agostino Gemelli. Fabi Alessandra reports a relationship with AIOM that includes: member of the national council. Meattini Icro reports a relationship with Menarini StemLine, Eli Lilly, AstraZeneca, Daiichi Sankyo, Novartis, Pfizer, Gilead, that includes: speaking and lecture fees.

Other authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors thank AIOM (Associazione Italiana di Oncologia Medica), Italy, for the open access funding. We would also like to thank "aBRCAadabra Onlus Association" for the valuable support in the consent review process, without which this effort would not have been possible.

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ctrv.2025.102976>.

References

- [1] Breast Cancer Association Consortium, Dorling L, Carvalho S, Allen J, González-Neira A, Luccarini C, Wahlström C, et al. Breast cancer risk genes - association analysis in more than 113,000 women. *N Engl J Med* 2021;384:428–39. <https://doi.org/10.1056/NEJMoa1913948>.
- [2] Garutti M, Foffano L, Mazzeo R, Michelotti A, Da Ros L, Viel A, et al. Hereditary cancer syndromes: a comprehensive review with a visual tool. *Genes (Basel)* 2023; 14(5):1025. <https://doi.org/10.3390/genes14051025>.
- [3] Fabi A, Cortesi L, Duranti S, Cordisco EL, Di Leone A, Terribile D, et al. Multigenic panels in breast cancer: clinical utility and management of patients with pathogenic variants other than BRCA1/2. *Crit Rev Oncol Hematol* 2024;201: 104431. <https://doi.org/10.1016/j.critrevonc.2024.104431>.
- [4] Mosele MF, Westphalen CB, Stenzinger A, Barlesi F, Bayle A, Bièche I, et al. Recommendations for the use of next-generation sequencing (NGS) for patients with advanced cancer in 2024: a report from the ESMO Precision Medicine Working Group. *Ann Oncol* 2024;35:588–606. <https://doi.org/10.1016/j.annonc.2024.04.005>.
- [5] Daly MB, Rosenthal E, Cummings S, Bernhisel R, Kidd J, Hughes E, et al. The association between age at breast cancer diagnosis and prevalence of pathogenic variants. *Breast Cancer Res Treat* 2023;199:617–26. <https://doi.org/10.1007/s10549-023-06946-8>.
- [6] Pujol P, Barberis M, Beer P, Friedman E, Piulats JM, Capoluongo ED, et al. Clinical practice guidelines for BRCA1 and BRCA2 genetic testing. *Eur J Cancer* 2021;146: 30–47. <https://doi.org/10.1016/j.ejca.2020.12.023>.
- [7] Hu C, Hart SN, Gnanaolivu R, Huang H, Lee KY, Na J, et al. A population-based study of genes previously implicated in breast cancer. *N Engl J Med* 2021;384: 440–51. <https://doi.org/10.1056/NEJMoa2005936>.
- [8] Fondazione AIOM-AIRTUM-PASSI. I Numeri del Cancro in Italia 2020. Available online: www.fondazioneaiom.it (Accessed February 9, 2025).
- [9] Tung NM, Zakalik D, Somerfield MR, Hereditary Breast Cancer Guideline Expert Panel. Adjuvant PARP inhibitors in patients with high-risk early-stage HER2-negative breast cancer and germline BRCA mutations: ASCO hereditary breast cancer guideline rapid recommendation update. *J Clin Oncol* 2021;39:2959–61. <https://doi.org/10.1200/JCO.21.01532>.
- [10] Tutt ANJ, Garber JE, Kaufman B, Viale G, Fumagalli D, Rastogi P, et al. Adjuvant olaparib for patients with BRCA1- or BRCA2-mutated breast cancer. *N Engl J Med* 2021;384:2394–405. <https://doi.org/10.1056/NEJMoa2105215>.
- [11] Robson M, Im SA, Senkus E, Xu B, Domchek SM, Masuda N, et al. Olaparib for metastatic breast cancer in patients with a germline BRCA mutation. *N Engl J Med* 2017;377:523–33. <https://doi.org/10.1056/NEJMoa1706450>.
- [12] Litton JK, Rugo HS, Ettl J, Hurvitz SA, Gonçalves A, Lee KH, et al. Talazoparib in patients with advanced breast cancer and a germline BRCA mutation. *N Engl J Med* 2018;379:753–63. <https://doi.org/10.1056/NEJMoa1802905>.
- [13] Schmid P, Cortes J, Pusztai L, McArthur H, Kümmel S, Bergh J, et al. Pembrolizumab for early triple-negative breast cancer. *N Engl J Med* 2020;382: 810–21. <https://doi.org/10.1056/NEJMoa1910549>.
- [14] Valachis A, Nearchou AD, Lind P. Surgical management of breast cancer in BRCA-mutation carriers: a systematic review and meta-analysis. *Breast Cancer Res Treat* 2014;144:443–55. <https://doi.org/10.1007/s10549-014-2890-1>.
- [15] Davey MG, Davey CM, Ryan EJ, Lowery AJ, Kerin MJ. Combined breast conservation therapy versus mastectomy for BRCA mutation carriers - a systematic review and meta-analysis. *Breast* 2021;56:26–34. <https://doi.org/10.1016/j.breast.2021.02.001>.
- [16] Pierce LJ, Phillips KA, Griffith KA, Buys S, Gaffney DK, Moran MS, et al. Local therapy in BRCA1 and BRCA2 mutation carriers with operable breast cancer: comparison of breast conservation and mastectomy. *Breast Cancer Res Treat* 2010; 121(2):389–98. <https://doi.org/10.1007/s10549-010-0894-z>.
- [17] Evron E, Ben-David AM, Goldberg H, Fried G, Kaufman B, Catane R, et al. Prophylactic irradiation to the contralateral breast for BRCA mutation carriers with early-stage breast cancer. *Ann Oncol* 2019;30:412–7. <https://doi.org/10.1093/annonc/mdy515>.
- [18] Kuchenbaecker KB, Hopper JL, Barnes DR, Phillips KA, Mooij TM, Roos-Blom MJ, et al. Risks of breast, ovarian, and contralateral breast cancer for BRCA1 and BRCA2 mutation carriers. *J Am Med Assoc* 2017;317:2402–16. <https://doi.org/10.1001/jama.2017.7112>.
- [19] Trainer AH, Lewis CR, Tucker K, Meiser B, Friedlander M, Ward RL. The role of BRCA mutation testing in determining breast cancer therapy. *Nat Rev Clin Oncol* 2010;7:708–17. <https://doi.org/10.1038/nrclinonc.2010.175>.
- [20] Cortesi L, Razzaboni E, Toss A, De Matteis E, Marchi I, Medici V, et al. A rapid genetic counselling and testing in newly diagnosed breast cancer is associated with high rate of risk-reducing mastectomy in BRCA1/2-positive Italian women. *Ann Oncol* 2014;25:57–63. <https://doi.org/10.1093/annonc/mdt422>.
- [21] Metcalfe KA, Lubinski J, Ghadirian P, Lynch H, Kim-Sing C, Friedman E, et al. Predictors of contralateral prophylactic mastectomy in women with a BRCA1 or BRCA2 mutation: the Hereditary Breast Cancer Clinical Study Group. *J Clin Oncol* 2008;26:1093–7. <https://doi.org/10.1200/JCO.2007.12.6078>.
- [22] Boughhey JC, Hoskin TL, Degnim AC, Sellers TA, Johnson JL, Kasner MJ, et al. Contralateral prophylactic mastectomy is associated with a survival advantage in high-risk women with a personal history of breast cancer. *Ann Surg Oncol* 2010;17: 2702–9. <https://doi.org/10.1245/s10434-010-1136-7>.
- [23] Bedrosian I, Hu CY, Chang GJ. Population-based study of contralateral prophylactic mastectomy and survival outcomes of breast cancer patients. *J Natl Cancer Inst* 2010;102:401–9. <https://doi.org/10.1093/jnci/djq018>.

- [24] Metcalfe KA, Lynch HT, Ghadirian P, Tung N, Olivetto IA, Foulkes WD, et al. The risk of ovarian cancer after breast cancer in BRCA1 and BRCA2 carriers. *Gynecol Oncol* 2005;96:222–6. <https://doi.org/10.1016/j.ygyno.2004.09.039>.
- [25] Kotsopoulos J, Gronwald J, Huzarski T, Møller P, Pal T, McCuaig JM, et al. Bilateral oophorectomy and all-cause mortality in women with BRCA1 and BRCA2 sequence variations. *JAMA Oncol* 2024;10(4):484–92. <https://doi.org/10.1001/jamaoncol.2023.6937>.
- [26] Ferlay J, Steliarova-Foucher E, Lortet-Tieulent J, Rosso S, Coebergh JW, Comber H, et al. Cancer incidence and mortality patterns in Europe: estimates for 40 countries in 2012. *Eur J Cancer* 2013;49:1374–403. <https://doi.org/10.1016/j.ejca.2012.12.027>.
- [27] Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics. *CA Cancer J Clin* 2012;2015:87–108. <https://doi.org/10.3322/caac.21262>.
- [28] Cortesi L, Cortesi G, Venturelli M, Marcheselli L, Toss A, Barbieri E, et al. Can contralateral prophylactic mastectomy and oophorectomy increase survival in BRCA-related breast cancer? Results from the Italian MUTina study. *Eur J Surg Oncol* 2024;50:108603. <https://doi.org/10.1016/j.ejso.2024.108603>.
- [29] Yadav S, Reeves A, Campian S, Sufka A, Zakalik D. Preoperative genetic testing impacts surgical decision making in BRCA mutation carriers with breast cancer: a retrospective cohort analysis. *Hered Cancer Clin Pract* 2017;15:11. <https://doi.org/10.1186/s13053-017-0071-z>.
- [30] Ain Q, Richardson C, Mutebi M, George A, Kemp Z, Rusby JE. Does mainstream BRCA testing affect surgical decision-making in newly-diagnosed breast cancer patients? *Breast* 2023;67:30–5. <https://doi.org/10.1016/j.breast.2022.12.001>.
- [31] Geyer Jr CE, Garber JE, Gelber RD, Yothers G, Taboada M, Ross L, et al. Overall survival in the OlympiA phase III trial of adjuvant olaparib in patients with germline pathogenic variants in BRCA1/2 and high-risk, early breast cancer. *Ann Oncol* 2022;33:1250–68. <https://doi.org/10.1016/j.annonc.2022.09.159>.
- [32] Schmid P, Cortes J, Dent R, McArthur H, Pusztai L, Kümmel S, et al. Overall survival with pembrolizumab in Early-stage triple-negative breast cancer. *N Engl J Med* 2024;391:1981–91. <https://doi.org/10.1056/NEJMoa2409932>.
- [33] Pusztai L, Denkert C, O'Shaughnessy J, Cortes J, Dent R, McArthur H, et al. Event-free survival by residual cancer burden with pembrolizumab in early-stage TNBC: exploratory analysis from KEYNOTE-522. *Ann Oncol* 2024;35:429–36. <https://doi.org/10.1016/j.annonc.2024.02.002>.
- [34] Safonov A, Marra A, Bandlamudi C, O'Leary B, Wubbenhorst B, Ferraro E, et al. Tumor suppressor heterozygosity and homologous recombination deficiency mediate resistance to front-line therapy in breast cancer. 2024.02.05.578934 bioRxiv [Preprint] 2024. <https://doi.org/10.1101/2024.02.05.578934>.
- [35] Meattini I, Becherini C, Caimi S, Coles CE, Cortes J, Curigliano G, et al. International multidisciplinary consensus on the integration of radiotherapy with new systemic treatments for breast cancer: European Society for Radiotherapy and Oncology (ESTRO)-endorsed recommendations. *Lancet Oncol* 2024;25(2):e73–83. [https://doi.org/10.1016/S1470-2045\(23\)00534-X](https://doi.org/10.1016/S1470-2045(23)00534-X).
- [36] Manahan ER, Kuerer HM, Sebastian M, Hughes KS, Boughey JC, Euhus DM, et al. Consensus guidelines on genetic testing for hereditary breast cancer from the American Society of Breast Surgeons. *Ann Surg Oncol* 2019;26:3025–31. <https://doi.org/10.1245/s10434-019-07549-8>.
- [37] Guo F, Adekanmbi V, Hsu CD, Berenson AB, Kuo YF, Shih YT. Cost-effectiveness of population-based multigene testing for breast and ovarian cancer prevention. *JAMA Netw Open* 2024;7:e2356078. <https://doi.org/10.1001/jamanetworkopen.2023.56078>.
- [38] Ferretti S, Sassoli de Bianchi P, Canuti D, Campari C, Cortesi L, Arcangeli V, et al. Evaluation of an Italian population-based programme for risk assessment and genetic counselling and testing for BRCA1/2-related hereditary breast and ovarian cancer after 10 years of operation: an observational study protocol. *Methods Protoc* 2024;7:63. <https://doi.org/10.3390/mps7040063>.
- [39] Cortesi L, Baldassarri B, Ferretti S, Razzaboni E, Bella M, Bucchi L, et al. A regional population-based hereditary breast cancer screening tool in Italy: first 5-year results. *Cancer Med* 2020;9:2579–89. <https://doi.org/10.1002/cam4.2824>.
- [40] Bokkers K, Bleiker EMA, Velthuisen ME, Koelmeij R, Burgmans JPI, Klinkenbijn JH, et al. Patients' experiences with pre-test genetic counseling provided by breast cancer healthcare professionals: results from a large prospective multicenter study. *Breast* 2023;69:349–57. <https://doi.org/10.1016/j.breast.2023.03.017>.
- [41] Burke K, Dawson L, Hodgkinson K, Wilson BJ, Etchegary H. Exploring family communication preferences in hereditary breast and ovarian cancer and lynch syndrome: a national Canadian survey. *J Commun Genet* 2024;15:387–400. <https://doi.org/10.1007/s12687-024-00720-z>.
- [42] Bono M, Fanale D, Incorvaia L, Cancelliere D, Fiorino A, Calò V. Impact of deleterious variants in other genes beyond BRCA1/2 detected in breast/ovarian and pancreatic cancer patients by NGS-based multi-gene panel testing: looking over the hedge. *ESMO Open* 2021;100235. <https://doi.org/10.1016/j.esmoop.2021.100235>.
- [43] Anacleiro F, Pilenzi L, Dell'Elice A, Ferrante R, Grossi S, Ferlito LM, et al. Clinical usefulness of NGS multi-gene panel testing in hereditary cancer analysis. *Front Genet* 2023;14:1060504. <https://doi.org/10.3389/fgene.2023.1060504>.
- [44] Tung NM, Robson ME, Ventz S, Santa-Maria CA, Nanda R, Marcom PK et al TBCRC 048: phase II study of Olaparib for metastatic breast cancer and mutations in homologous recombination-related genes. *J Clin Oncol* 2020;38:4274–82. <https://doi.org/10.1200/JCO.20.02151>.
- [45] Makhnoon S, Shirts BH, Bowen DJ. Patients' perspectives of variants of uncertain significance and strategies for uncertainty management. *J Genet Couns* 2019;28:313–25. <https://doi.org/10.1002/jgc4.1075>.
- [46] O'Neill SC, Rini C, Goldsmith RE, Valdimarsdottir H, Cohen LH, Schwartz MD. Distress among women receiving uninformative BRCA1/2 results: 12-month outcomes. *Psychooncology* 2009;18:1088–96. <https://doi.org/10.1002/pon.1467>.
- [47] Scherr CL, Ramesh S, Getachew-Smith H, Kalke K, Ramsey K, Fischhoff B, et al. How patients deal with an ambiguous medical test: decision-making after genetic testing. *Patient Educ Couns* 2021;104:953–9. <https://doi.org/10.1016/j.pec.2020.10.020>.
- [48] Makhnoon S, Bednar EM, Krause KJ, Peterson SK, Lopez-Olivo MA. Clinical management among individuals with variant of uncertain significance in hereditary cancer: a systematic review and meta-analysis. *Clin Genet* 2021;100(2):119–31. <https://doi.org/10.1111/cge.13966>.
- [49] Ro V, McGuinness JE, Guo B, Trivedi MS, Jones T, Chung WK, et al. Association between genetic testing for hereditary breast cancer and contralateral prophylactic mastectomy among multiethnic women diagnosed with early-stage breast cancer. *JCO Oncol Pract* 2022;18(4):e472–83. <https://doi.org/10.1200/OP.21.00322>.
- [50] Makhnoon S, Chen M, Levin B, Ensinger M, Mattie KD, Grana G, et al. Use of breast surveillance between women with pathogenic variants and variants of uncertain significance in breast cancer susceptibility genes. *Cancer* 2022;128(20):3709–17. <https://doi.org/10.1002/cncr.34429>.
- [51] Fridman AL, Raz A, Timmermans S, Shkedi-Rafid S. Fragmented responsibility: views of Israeli HCPs regarding patient recontact following variant reclassification. *J Commun Genet* 2022;13:13–8. <https://doi.org/10.1007/s12687-021-00556-x>.
- [52] Kobayashi Y, Chen E, Facio FM, Metz H, Poll SR, Swartzlander D, et al. Clinical variant reclassification in hereditary disease genetic testing. *JAMA Netw Open* 2024;7:e2444526. <https://doi.org/10.1001/jamanetworkopen.2024>.
- [53] Bedrosian I, Somerfield MR, Achatz MI, Boughey JC, Curigliano G, Friedman S, et al. Germline testing in patients with breast cancer: ASCO-society of surgical oncology guideline. *J Clin Oncol* 2024;42:584–604. <https://doi.org/10.1200/JCO.23.02225>.
- [54] Kurian AW, Bernhisel R, Larson K, Caswell-Jin JL, Shadyab AH, Ochs-Balcom H, et al. Prevalence of pathogenic variants in cancer susceptibility genes among women with postmenopausal breast cancer. *J Am Med Assoc* 2020;323:995–7. <https://doi.org/10.1001/jama.2020.0229>.
- [55] Lazard T, Bataillon G, Naylor P, Popova T, Bidard FC, Stoppa-Lyonnet D, et al. Deep learning identifies morphological patterns of homologous recombination deficiency in luminal breast cancers from whole slide images. *Cell Rep Med* 2022;3:100872. <https://doi.org/10.1016/j.xcrm.2022.100872>.
- [56] Bergstrom EN, Abbasi A, Díaz-Gay M, Galland L, Ladoire S, Lippman SM, et al. Deep learning artificial intelligence predicts homologous recombination deficiency and platinum response from histologic slides. *J Clin Oncol* 2024;42:3550–60. <https://doi.org/10.1200/JCO.23.02641>.