

Rafoxanide negatively modulates STAT3 and NF- κ B activity and inflammation-associated colon tumorigenesis

Teresa Pacifico¹ | Carmine Stolfi¹ | Lorenzo Tomassini¹ | Anderson Luiz-Ferreira² | Eleonora Franzè¹ | Angela Ortenzi¹ | Alfredo Colantoni¹ | Giuseppe S. Sica³ | Manolo Sambucci⁴ | Ivan Monteleone⁵ | Giovanni Monteleone¹ | Federica Laudisi¹ 

¹Department of Systems Medicine, University of Rome "Tor Vergata", Rome, Italy

²Inflammatory Bowel Disease Research Laboratory, Department of Biological Sciences, Institute of Biotechnology, Federal University of Catalão (UFCAT), Catalão, Brazil

³Department of Surgery, University of Rome "Tor Vergata", Rome, Italy

⁴Neuroimmunology Unit, Santa Lucia Foundation IRCCS, Rome, Italy

⁵Department of Biomedicine and Prevention, University of Rome "Tor Vergata", Rome, Italy

Correspondence

Federica Laudisi, Department of Systems Medicine, University of Rome "Tor Vergata", Via Montpellier, 1, 00133 Rome, Italy.
Email: federica.laudisi@uniroma2.it

Funding information

Fondazione AIRC per la Ricerca sul Cancro ETS, Grant/Award Number: MFAG-27142; Ministero dell'Università e della Ricerca (MUR), Grant/Award Number: PNRR-M4C2I1.3PE6 and 2022JEBP88

Abstract

In the colorectal cancer (CRC) niche, the transcription factors signal transducer and activator of transcription 3 (STAT3) and nuclear factor- κ B (NF- κ B) are hyperactivated in both malignant cells and tumor-infiltrating leukocytes (TILs) and cooperate to maintain cancer cell proliferation/survival and drive protumor inflammation. Through drug repositioning studies, the anthelmintic drug rafoxanide has recently emerged as a potent and selective antitumor molecule for different types of cancer, including CRC. Here, we investigate whether rafoxanide could negatively modulate STAT3/NF- κ B and inflammation-associated CRC. The antineoplastic effect of rafoxanide was explored in a murine model of CRC resembling colitis-associated disease. Cell proliferation and/or STAT3/NF- κ B activation were evaluated in colon tissues taken from mice with colitis-associated CRC, human CRC cells, and CRC patient-derived explants and organoids after treatment with rafoxanide. The STAT3/NF- κ B activation and cytokine production/secretion were assessed in TILs isolated from CRC specimens and treated with rafoxanide. Finally, we investigated the effects of TIL-derived supernatants cultured with or without rafoxanide on CRC cell proliferation and STAT3/NF- κ B activation. The results showed that rafoxanide restrains STAT3/NF- κ B activation and inflammation-associated colon tumorigenesis in vivo without apparent effects on normal intestinal cells. Rafoxanide markedly reduces STAT3/NF- κ B activation in cultured CRC cells, CRC-derived explants/organoids, and TILs. Finally, rafoxanide treatment impairs the ability of TILs to produce protumor cytokines and promote CRC cell proliferation. We report the novel observation that rafoxanide negatively affects STAT3/NF- κ B oncogenic activity at multiple levels in the CRC microenvironment. Our data suggest that rafoxanide could potentially be deployed as an anticancer drug in inflammation-associated CRC.

KEYWORDS

anthelmintic drug, colitis-associated colorectal cancer, cytokine, drug repositioning, tumor microenvironment

Teresa Pacifico and Carmine Stolfi contributed equally to this work.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2024 The Author(s). *Cancer Science* published by John Wiley & Sons Australia, Ltd on behalf of Japanese Cancer Association.

1 | INTRODUCTION

Chronic inflammation is recognized as an important player in the pathogenesis of many types of neoplasia, as it can promote the onset, progression, and invasion of cancer cells.¹ In particular, in patients with ulcerative colitis, one of the main forms of chronic inflammatory bowel disease in humans, there is an increased risk of developing colorectal cancer (CRC),² which is related to the duration, extent, and severity of the inflammatory condition.^{3,4} However, it is worth mentioning that even sporadic CRCs, which represent the majority of CRC cases, exhibit extensive immune cell infiltrates and prominent inflammatory responses.^{5,6}

The transcription factors signal transducer and activator of transcription 3 (STAT3) and nuclear factor- κ B (NF- κ B) lie at the hub of multiple signaling pathways, playing a central role in a multitude of physiological processes, including inflammation, immune cell development, cell proliferation, and survival.^{7,8} Typically, STAT3 activation is triggered by binding of cytokines and growth factors to their related receptors. For example, members of the interleukin-6 (IL-6) family of cytokines are potent STAT3 activators, mediating their signal through the pleiotropic gp130 receptor subunit that recruits and triggers the activation of JAK2 by transphosphorylation.⁹ JAK2 can then phosphorylate STAT3 on the Y705 residue.¹⁰ Phosphorylated STAT3 can homodimerize, as well as form dimers with STAT1, and then move into the nucleus to exert its functions.¹¹

Nuclear factor- κ B consists of a family of related transcription factors (i.e., RelA/p65, RelB, c-Rel, p50, and p52) that can hetero- and homodimerize to form at least 12 different identified dimers, and whose interactions, stability, degradation, and transcription activity are controlled by site-specific phosphorylation/s of NF- κ B subunits within transactivation domains.¹² In particular, p65 Serine 536 (S536) phosphorylation¹³ has been reported to promote its nuclear retention, enhance its association with transcriptional coactivators, inhibit its association with corepressors, and potentiate the activation of NF- κ B target genes.¹³⁻¹⁷ Dysregulation of STAT3/NF- κ B contributes to the development of various autoimmune and inflammatory diseases,¹⁸⁻²⁰ as well as malignant disorders.²¹ In this latter context, accumulating evidence produced in different types of CRC indicates that STAT3 and NF- κ B are constitutively activated in both malignant cells and tumor-infiltrating leukocytes (TILs), playing a key role in maintaining cancer cell proliferation/survival, mediating the interaction between cancer cells and immune/inflammatory cells, upregulating the release of inflammatory mediators (i.e., IL-6, tumor necrosis factor- α [TNF- α], and IL-1 β cytokines), triggering tumor angiogenesis and invasiveness.^{19,21,22} Given the pleiotropic effects of these two transcription factors in the CRC niche, compounds inhibiting STAT3 and NF- κ B inappropriate/persistent activation represent promising therapeutic tools and could help combat this neoplasia.

By drug repositioning studies – defined as the investigation of existing drugs for the treatment of pathologies that fall outside the scope of the original medical indication – the halogenated salicylanilide compound rafoxanide, an anthelmintic drug commonly used for the veterinary treatment of fascioliasis and some gastrointestinal

roundworms, has recently emerged as a potent antitumor agent for different types of cancer.²³⁻²⁶ In particular, we reported that rafoxanide exerted selective antimetagenic and proapoptotic effects on CRC cells in vitro and in an experimental model mimicking sporadic CRC.²⁷ More recently, our follow-up studies in cultured cells and preclinical CRC models identified rafoxanide as a bona fide immunogenic cell death inducer and a selective sensitizer to TNF-related apoptosis-inducing ligand-driven cell death.^{28,29}

In this study, we aimed to investigate whether rafoxanide could negatively modulate STAT3/NF- κ B activity and inflammation-associated colon tumorigenesis.

2 | MATERIALS AND METHODS

2.1 | Patients

Tissue samples were taken from the tumor area of 12 patients who underwent colon resection for sporadic CRC (all with TNM stages II–III) at the Tor Vergata University Hospital and used for either the generation of organoids or the isolation of TILs. No patients received radiation therapy or chemotherapy prior to surgery. Human studies were approved by the local ethics committee (protocol no. R.S. 131.17) and each patient gave written informed consent. The study methodologies were in accordance with the standards set by the Declaration of Helsinki.

2.2 | Mice

C57BL/6J female mice (6–8 weeks of age) were purchased from Charles River Laboratories Italia Srl. The mice were housed in standard animal cages under specific pathogen-free conditions in the animal facility of the University of Rome “Tor Vergata”. Mice were routinely tested (every 3 months) for health status and infections according to the guidelines of the Federation of European Laboratory Animal Science Associations (FELASA). Mice tested negative for all pathogens were included in this protocol. All in vivo experiments were approved by the animal ethics committee according to the Italian legislation on animal experimentation (authorization no. 494/2017-PR) and in accordance with European rules (2010/63/UE).

2.3 | Cell cultures

Unless otherwise noted, reagents were purchased from Merck Life Science. The human CRC cell line HCT-116 was obtained from ATCC and was maintained in McCoy's 5A medium supplemented with 10% FBS and 1% penicillin/streptomycin (P/S). The human CRC cell line DLD1 was obtained from ATCC and was maintained in RPMI-1640 medium supplemented with 10% FBS and 1% P/S. Cell lines were recently authenticated by short tandem repeat

(STR) DNA fingerprinting using the PowerPlex 18D System kit according to the manufacturer's instructions (Promega). The STR profiles of all cell lines matched the known DNA fingerprints. Rafoxanide (PubChem CID: 31475) was purchased from Tokyo Chemical Industry Co., Ltd.

To test whether rafoxanide inhibited STAT3 and/or NF- κ B activation in CRC cells, HCT-116 and DLD1 cells were incubated with increasing doses of the drug (1.25–5 μ M) for 24 h or with 2.5 μ M rafoxanide for 1–6 h.

To shed light on the mechanism/s by which rafoxanide modulates STAT3 Y705 phosphorylation, HCT-116 and DLD1 cells were preincubated or not with the general protein tyrosine phosphatase (PTP) inhibitor Na₃VO₄ (used at 100 μ M) for 1 h, and then stimulated with DMSO or 2.5 μ M rafoxanide for further 24 h.

To assess the effect of supernatants derived from TILs cultured in the presence of DMSO (TIL SN) or rafoxanide (RFX TIL SN), HCT-116 and DLD1 cells were cultured in the presence of TIL SN, RFX TIL SN, or RPMI-1640 medium (vehicle) (all used at 1:20 final dilution). Cell proliferation was assessed after 24 h using a BrdU assay (Roche Diagnostics) and the expression of phosphorylated (p)-STAT3 Y705 and p-NF- κ B/p65 S536 was evaluated after 30 min by western blotting.

2.4 | Experimental models of colitis-associated and sporadic CRC

To induce colitis-associated CRC (CAC), cohoused 6–8-week-old female C57BL/6J mice received a first i.p. injection of azoxymethane (AOM; 10 mg/kg) on day 0. After 7 days, mice were given 2% dextran sulfate sodium (DSS) (molecular weight, 9000–20,000; MP Biomedicals LLC) dissolved in the drinking water for 4 consecutive days to induce colitis. Mice received a second i.p. injection of AOM on day 18 (5 mg/kg) and were again given 2% DSS in the drinking water on day 25 for 4 days. The mice were then given regular water until the end of the experiment (day 90). On day 36 mice were randomly divided into two groups and given 7.5 mg/kg rafoxanide (in 10% DMSO in PBS) or 10% DMSO in PBS (control) every other day by i.p. injection until they were killed (day 90). Bodyweight was recorded every week from day 36 to the end of the study. Colonoscopy was carried out on day 88 in a blinded manner for monitoring of tumorigenesis using the Coloview high-resolution mouse endoscopic system (Karl Storz). The lesions observed during endoscopy were counted to obtain the total number of lesions. The size of all lesions in a given mouse was scored using the protocol described by Becker et al.³⁰ Cohoused 6–7-week-old female *Apc*^{min/+} mice received i.p. injections of 10 mg/kg AOM once a week for 2 weeks to increase colon tumorigenesis as previously reported.⁶ Two weeks after the last injection of AOM, mice were randomly divided into two groups and given either 7.5 mg/kg rafoxanide (in 10% DMSO in PBS) or 10% DMSO in PBS (control) every other day by i.p. injections until they were killed (day 90). The dose of rafoxanide was selected according to that currently used in veterinary treatment (i.e., 7.5–10 mg/kg).

2.5 | Western blot analysis

Total proteins were extracted from colon tissue samples isolated from C57BL/6J and *Apc*^{min/+} mice, from CRC cell lines, and TILs using the following lysis buffer: 10 mmol/L HEPES, 1 mmol/L EDTA pH 8.0, 60 mmol/L KCl, 0.2% IGEPAL CA-630 (Nonidet P-40), 1 mmol/L sodium fluoride, 10 μ g/mL aprotinin, 10 μ g/mL leupeptin, 1 mmol/L DTT, 1 mmol/L PMSF, and 1 mmol/L sodium orthovanadate, separated on an SDS-PAGE gel and transferred using a Trans-Blot Turbo apparatus (Bio-Rad Laboratories). The samples were then incubated with the following Abs: p-STAT3 Y705 (#9145), p-NF- κ B/p65 S536 (#3033), IKK α (#61294), and IKK β (#8943) (1:1000 final dilution; Cell Signaling Technology), STAT3 (sc-8019), NF- κ B/p65 (sc-372), SHP-1 (sc-7289), SHP-2 (sc-7384), JAK2 (sc-390,539), and CDK6 (sc-177) (1:500 final dilution; Santa Cruz Biotechnology, Inc.) and β -actin (A-1978; 1:5000 final dilution) followed by a secondary Ab conjugated to HRP (1:20,000; Dako). Membrane acquisition was carried out in chemiluminescence with the ChemiDoc Imaging System (Bio-Rad Laboratories). Computer-assisted scanning densitometry (Image-Lab 5.2.1; Bio-Rad Laboratories) was used to analyze the intensity of the immunoreactive bands.

2.6 | Enzyme-linked immunosorbent assay

Total proteins extracted from colon tissue samples isolated from C57BL/6J and *Apc*^{min/+} mice were evaluated for the presence of IL-6 and TNF- α by specific ELISA kits (M600OB and MTA00B, respectively; Bio-Techne s.r.l.) according to the manufacturer's instructions. The levels of IL-6 and TNF- α in TIL SN and RFX TIL SN were evaluated using specific ELISA kits (EHL-IL6 and EHL-TNF α , respectively; RayBiotech) according to the manufacturer's instructions.

2.7 | Organ culture

Organ culture experiments were carried out as previously described.³¹ Briefly, human CRC explants were placed on Millicell inserts (Min a 6-well plate containing complete RPMI-1640 medium supplemented with 50 μ g/mL gentamycin in the presence of DMSO (vehicle) or 2.5 μ M rafoxanide for 12 h. Culture was carried out in an organ culture chamber at 37°C in a 5% CO₂/95% O₂ atmosphere.

2.8 | Generation and culture of patient-derived CRC organoids

The resected intestinal tumor tissues were rinsed in HBSS containing antibiotics and cut into 5 mm pieces. These tissue fragments were incubated in Advanced DMEM/F12 medium (Thermo Fisher Scientific) containing 15 mM EDTA and rocked at 4°C for 30 min. The larger tissue fragments were removed and the remaining crypts were centrifugated at 300g for 5 min, embedded in Matrigel (CLS356231), and

plated in a warmed 24-well plate. The Matrigel was allowed to solidify for 30 min at 37°C and overlaid with complete medium (#06010 IntestiCult OGM Human; StemCell Technologies) supplemented with 10 μ M Y27632, a selective inhibitor of Rho-associated, coiled-coil containing protein kinase (ROCK) (#72302; StemCell Technologies). The entire medium was replaced every 3 days. The established organoids were then cultured with either DMSO or rafoxanide (2.5 μ M) for 24 h. The culture medium was then removed and the organoids were washed with PBS and incubated with organoid harvesting solution (3700-100-01; Bio-Techne s.r.l.) for 1 h at 4°C with gentle shaking. The released organoids from the depolymerized Matrigel were then collected and transferred to a cryomold containing optimal cutting temperature compound, frozen, and stored at -80°C.

2.9 | Immunohistochemistry

Cryosections of colon tissue samples isolated from C57BL/6J undergoing AOM/DSS were stained with a primary Ab directed against Ki-67 (clone TEC-3, M7249; Dako), p-eukaryotic translation initiation factor 2 α (eIF2 α) (Ser51; #3597), and cleaved caspase-3 (Asp175; #9661) (both from Cell Signaling Technology). Cryosections of human CRC explants and CRC patient-derived organoids were stained with a primary Ab against p-STAT3 Y705 (#9145; Cell Signaling Technology), p-NF- κ B/p65 S536 (#3033; Cell Signaling Technology), or Ki-67 (clone MIB-1 sc-101,861; Santa Cruz Biotechnology). Positive cells were visualized using the MACH4 Universal HRP Polymer kit with DAB (Biocare Medical) and analyzed using the Leica DMI4000 B microscope using the Leica application suite software (version 4.6.2).

2.10 | Isolation and culture of TILs

Tumor-infiltrating leukocytes were isolated from the lesions of AOM/DSS treated mice by enzymatic digestion with Liberase TM (200 μ g/mL, 05401127001; Roche Diagnostics) and DNase I (200 μ g/mL, 11284932001; Roche Diagnostics). Cells were filtered through a 70 μ m cell strainer, washed in HBSS, and resuspended in complete RPMI-1640 medium.

To isolate human TILs, tumor tissue pieces were dissected from surgical samples within 1 h after resection and rinsed in HBSS containing antibiotics. The samples were then incubated in HBSS containing 1 mM EDTA and antibiotics for 45 min at 37°C to remove epithelial cells. After two washes in HBSS, the samples were minced and incubated with Liberase TM (200 μ g/mL, 05401127001; Roche Diagnostics) and DNase I (200 μ g/mL, 11284932001; Roche Diagnostics) for 1 h at 37°C. After Liberase digestion, media containing mononuclear cells were collected and washed twice in HBSS. Subsequently, the pellets were resuspended in complete RPMI-1640 and then layered on a Percoll density gradient to isolate the TILs. After two washes in complete RPMI-1640 medium, isolated cells were counted and viability was verified using 0.1% Trypan blue (viability ranged from 90% to

97%). Freshly isolated TILs were resuspended in complete RPMI-1640 medium. An aliquot of cells was seeded and stimulated with DMSO (vehicle) or rafoxanide (2.5 μ M) for 16 h, cells were then collected and STAT3/NF- κ B activation was assessed by western blotting. An aliquot of cells was cultured in 96-well plates in the presence of DMSO or rafoxanide (2.5 μ M) for 16 h. Phorbol 12-myristate 13-acetate (80 pM), ionomycin (1 mg/mL), and monensin (2 μ M; eBioscience) were added to the cultures 4 h before the end of stimulation. Cells were then collected and the fraction of IL-6 and/or TNF- α producing TILs were assessed by flow cytometry after staining with specific fluorochrome-conjugated Abs. One million TILs were cultured in 48-well plates in the presence of DMSO or rafoxanide (2.5 μ M) to produce 0.5 mL supernatant. Cell-free supernatants were harvested after 16 h.

2.11 | Flow cytometry

Human TILs were stained with the following reagents/mAbs: LIVE/DEAD Fixable Violet Dead Cell Stain kit (L34955; Thermo Fisher Scientific), CD45-eFluor 506 (69-0459-42; eBioscience), CD3-APC-H7 (560176; BD Biosciences), γ -interferon (IFN- γ)-FITC (554700; BD Biosciences), IL-6-PerCP (46-7069-42; eBioscience), and TNF- α -PE (12-7349-82; eBioscience). In parallel, cells were stained with the appropriate control isotype Abs. Mouse TILs were stained with the following reagents/mAbs: LIVE/DEAD Fixable Aqua Dead Cell Stain kit (L34966; Thermo Fisher Scientific), CD45 APC-Cy7 (557659; BD Biosciences), CD3-Pacific Blue (558214; BD Biosciences), CD8-PerCP (45008182; BD Biosciences), DX5-PE (553858; BD Biosciences), IFN- γ FITC (554411; BD Biosciences), Perforin APC (17-9392-80; eBioscience), and granzyme B-Alexa Fluor 700 (372222; BioLegend). In parallel, cells were stained with the appropriate control isotype Abs. Flow cytometry analysis was undertaken using a Gallios flow cytometer (Beckman Coulter).

2.12 | Statistical analysis

Parametric data were analyzed using the two-tailed Student's *t*-test for comparison between two groups or one-way ANOVA followed by Tukey's or Dunnett's post hoc tests for multiple comparisons. Significance was defined as $p < 0.05$.

3 | RESULTS

3.1 | Rafoxanide reduces neoplastic cell proliferation/survival and STAT3/NF- κ B activation in a mouse model of CAC

In initial studies, we evaluated the possible antineoplastic effect of rafoxanide in a mouse model mimicking human CAC whereby AOM administration followed by repeated DSS ingestion caused colonic

inflammation and the subsequent development of multiple colonic lesions. To rule out the possibility that rafoxanide could interfere with the ongoing mucosal inflammation, drug treatment was started 1 week after discontinuation of the last treatment with DSS (Figure 1A). At that time (day 36), no visible lesions were detected by endoscopy (not

shown). Mice received i.p. injections of rafoxanide (7.5 mg/kg/mouse) or vehicle (control) every other day, until the mice were killed at the end of the study (Figure 1A). Endoscopy on day 88 showed that control mice developed multiple large colonic lesions, whereas the number and size of lesions were reduced in the colons of mice treated




FIGURE 1 Rafoxanide (RFX) limits colonic tumorigenesis in the azoxymethane (AOM)/dextran sulfate sodium (DSS)-driven colitis-associated colorectal cancer (CAC) model. (A) Schematic overview of the protocol used to induce CAC in mice. (B) Left panels show representative endoscopic pictures of colon lesions developed in mice treated with AOM/DSS and receiving vehicle (CTR) or RFX. Graphs show the number of lesions (tumor score) and the endoscopic score (tumor load) of the lesions developed in mice treated as indicated above. Data indicate the mean $\pm$ SEM of three independent experiments in which at least four mice per group were considered. (C) Representative images showing Ki-67 immunostaining in colonic sections taken from mice treated as indicated in (B). Scale bars, 20 μ m; 10 μ m (insets). One of five representative experiments in which similar results were obtained is shown. Right inset, quantification of Ki-67⁺ epithelial cells in colonic sections taken from mice treated as indicated in (B). Data are presented as mean values of positive cells per high power field $\pm$ SEM of two independent experiments in which at least two sections per group were analyzed. NT, nontumor area; T, tumor area.

with rafoxanide (Figure 1B). No significant changes in bodyweight were observed in mice treated with rafoxanide compared to control mice (not shown), in line with rafoxanide treatment experiments performed in healthy mice (pers. obs., 2024). Immunohistochemistry for Ki-67 (Figure 1C) and cleaved caspase-3 (Figure S1A) confirmed the antimitogenic and proapoptotic effects of rafoxanide on neoplastic cells. It should be noted that no significant change in Ki-67 or cleaved caspase-3 staining (Figures 1C and S1A) was detected in the normal colonic mucosa of mice treated with the drug, according to previously reported observations in a mouse model mimicking sporadic CRC.²⁷ In the AOM/DSS-induced lesions of rafoxanide-treated mice, the increase in the phosphorylation of eIF2 α on the S51 residue in transformed colon epithelial cells (Figure S1B), a mandatory event for cells undergoing immunogenic death to emit all the signals required for the initiation of the host immune response,³² as well as the expanded fraction of T lymphocytes and NK cells producing IFN- γ , a key cytokine in the activation of cellular immunity and, subsequently, stimulation of antitumor immune response,³³ and perforin (Figure S2), are in line with our previous report indicating rafoxanide as a bona fide immunogenic cell death inducer.²⁸

Excessive activation of STAT3 and NF- κ B transcription factors makes a major contribution to the tumorigenic process in the colon.²¹ To determine whether the reduced carcinogenesis observed in mice treated with rafoxanide was associated with a reduced activation of STAT3 and/or NF- κ B, we compared the expression of p-STAT3 Y705 and p-NF- κ B/p65 S536 in nontumor and tumor tissues derived from colon extracts of

control and rafoxanide-treated mice killed on day 90. Robust activation of both STAT3 and NF- κ B was observed in the neoplastic areas of control mice compared to those derived from mice treated with rafoxanide (Figure 2A). In the microenvironment characterizing colitis-associated colorectal carcinogenesis, in addition to intrinsic molecular defects in the signaling machinery, aberrant activation of STAT3 and/or NF- κ B that occurs in transformed epithelial cells can be increased by the presence of immune cell-derived protumorigenic cytokines.⁵ Therefore, next we assessed the expression of IL-6 and TNF- α , among the most potent activators of STAT3 and NF- κ B in colonic epithelial cells, respectively,^{5,34} in nontumor and tumor colonic tissues taken from mice treated as indicated above. Both IL-6 and TNF- α were reduced in tumors of mice that underwent drug treatment compared to control mice, while such cytokines were barely detectable in nontumor areas of control and rafoxanide-treated mice (Figure 2B). Taken together, our data indicate that rafoxanide reduces the development of neoplastic lesions driven by AOM/DSS colitis and this effect is associated with reduced STAT3/NF- κ B activation and production of IL-6 and TNF- α .

3.2 | Rafoxanide decreases STAT3/NF- κ B activation in colonic neoplastic lesions arising in APC^{min/-} mice and in human CRC explants

In addition to cancers of the digestive tract arising on overt inflammation, sporadic CRC, which represents the majority of CRC cases,

FIGURE 2 Reduced signal transducer and activator of transcription 3 (STAT3)/nuclear factor- κ B (NF- κ B) activation and interleukin-6 (IL-6)/tumor necrosis factor- α (TNF- α) expression in lesions of mice treated with azoxymethane (AOM)/dextran sulfate sodium (DSS) receiving rafoxanide (RFX). (A) Colonic extracts from mice treated with AOM/DSS receiving vehicle (CTR) or RFX and killed on day 90 were analyzed for phosphorylated (p-)STAT3 Y705, STAT3, p-NF- κ B/p65 S536, and NF- κ B/p65 expression by western blotting. β -Actin was used as a loading control. One of four representative experiments in which similar results were obtained is shown. (B) Scatter plots showing IL-6 and TNF- α protein expression assessed by ELISA in colonic extracts of mice treated as indicated in (A). Values are mean $\pm$ SEM of four experiments. NT, nontumor area; T, tumor area.

