

REVIEW

WILEY

Procedure-related bleeding risk in patients with cirrhosis and severe thrombocytopenia

Domenico Alvaro¹ | Nicola Caporaso² | Edoardo Giovanni Giannini³  | Angelo Iacobellis⁴ | Mariacristina Morelli⁵ | Pierluigi Toniutto⁶ | Francesco Violi⁷ | The PREBRiC (Procedure-Related Bleeding Risk in Cirrhosis) group

¹Department of Translational and Precision Medicine, Sapienza University of Rome, Rome, Italy

²Department of Clinical Medicine and Surgery, University of Naples 'Federico II', Naples, Italy

³Gastroenterology Unit, Department of Internal Medicine, University of Genoa, IRCCS-Ospedale Policlinico San Martino, Genoa, Italy

⁴Division of Gastroenterology, Fondazione IRCCS Casa Solievo della Sofferenza, Foggia, Italy

⁵IRCCS Azienda Ospedaliero-Universitaria di Bologna, Bologna, Italy

⁶Hepatology and Liver Transplantation Unit, Azienda Sanitaria Universitaria Integrata, Academic Hospital, Udine, Italy

⁷I Clinica Medica, Sapienza University of Rome, Rome, Italy

Correspondence

Edoardo Giovanni Giannini,
Gastroenterology Unit, Department of Internal Medicine, University of Genoa, IRCCS-Ospedale Policlinico San Martino, Viale Benedetto XV, no. 6 – 16132, Genoa, Italy.
Email: egiannini@unige.it

Funding information

Shionogi srl

Abstract

Background: Gaps of knowledge still exist about the potential association between severe thrombocytopenia and increased risk of procedure-associated bleeding in patients with liver disease.

Methods: In this narrative review, we aimed at examining the association between procedure-related bleeding risk and platelet count in patients with cirrhosis and severe thrombocytopenia in various settings. We updated to 2020 a previously conducted literature search using MEDLINE/PubMed and EMBASE. The search string included clinical studies, adult patients with chronic liver disease and thrombocytopenia undergoing invasive procedures, any interventions and comparators, and haemorrhagic events of any severity as outcome.

Results: The literature search identified 1276 unique publications, and 15 studies met the inclusion criteria and were analysed together with those identified by the previous search. Most of the new studies included in our analysis did not assess the association between post-procedural bleeding risk and platelet count alone in patients with chronic liver disease. Furthermore, some results could have been biased by prophylactic platelet transfusions. A few studies found that severe thrombocytopenia may be predictive of bleeding following percutaneous liver biopsy, dental extractions, percutaneous ablation of liver tumours and endoscopic polypectomy.

Conclusions: Currently available literature cannot support definitive conclusions about the appropriate target platelet counts to improve the risk of bleeding in cirrhotic patients who underwent invasive procedures; moreover, it showed enormous variability in the use of prophylactic platelet transfusions.

KEYWORDS

biopsy, liver cirrhosis, platelet count, platelet transfusions, thrombocytopenia

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. European Journal of Clinical Investigation published by John Wiley & Sons Ltd on behalf of Stichting European Society for Clinical Investigation Journal Foundation.

1 | INTRODUCTION

Thrombocytopenia is a very common complication of chronic liver disease.¹ However, severity of the underlying liver disease and differences in definition criteria of low platelet count cut-off makes the prevalence of thrombocytopenia widely variable.^{2,3} Available data show a prevalence of thrombocytopenia (i.e. platelet count $< 150 \times 10^9/L$) ranging from 6% to 78%, with lower percentages in non-cirrhotic patients, which become progressively greater in patients with compensated and decompensated cirrhosis.^{1,4,5} Also, moderate (i.e. platelet count between 50 and $75 \times 10^9/L$) and severe ($< 50 \times 10^9/L$) thrombocytopenia has been reported in 13% and 1% of patients with cirrhosis, respectively.⁶

Despite the possible coexistence of coagulopathy, mild-to-moderate thrombocytopenia (i.e. platelet count between 50 and $150 \times 10^9/L$) rarely represents a critical condition in patients without complication of liver disease (e.g. infections and renal failure). On the contrary, a platelet count $< 50 \times 10^9/L$ could have a negative impact on the clinical management of patients with advanced liver disease, since it may lead to postponement or cancellation of invasive procedures and may be associated with an increased risk of procedure-associated bleeding.⁶⁻⁸

De Pietri et al⁹ categorized procedures based on the associated bleeding risk: procedures were defined at high or low risk if associated with a bleeding risk $> 3\%$ (variceal band ligation, hepatic resection, abdominal surgery, endoscopic polypectomy, radio-frequency ablation, liver biopsy, biopsy of sites other than liver, abdominal drainage and endoscopic retrograde cholangiopancreatography with sphincterotomy or thoracotomy) or $< 3\%$ (paracentesis, thoracentesis, central vein cannulation and transjugular intrahepatic portosystemic shunts), respectively.

Literature analysis conducted by the working group of the Position Paper of the Italian Association for the Study of Liver Diseases (AISF) and the Italian Society of Internal Medicine (SIMI)⁸ found that the bleeding risk in cirrhotic patients was low (3%) following paracentesis, thoracentesis, and percutaneous or transjugular liver biopsy, and moderate ($< 10\%$) following endoscopic variceal ligation, endoscopic polypectomy and minor abdominal surgery (i.e. cholecystectomy and hernioplasty). The Position Paper concluded that, despite the limitations of the studies analysed, a platelet count $< 50-60 \times 10^9/L$ may be predictive of procedure-associated bleeding.⁸

The aim of this narrative review was to update the literature search conducted by the AISF/SIMI working group and examine the association between procedure-related bleeding risk and the platelet count in patients with cirrhosis and severe thrombocytopenia in different settings.

2 | LITERATURE SEARCH

Starting from the literature search conducted by the working group of the Position Paper AISF/SIMI, which covered relevant evidence on 'Risk of bleeding following invasive procedures or surgery' until 2014,⁸ we conducted a new search using MEDLINE/PubMed and EMBASE with the aim to update data from 2014 to 2020 (last accessed on 4 March 2020). The search string has been designed on the basis of the PICOS scheme and included clinical (RCT and observational) studies conducted on adult patients with chronic liver disease and thrombocytopenia undergoing invasive procedures, with any interventions and comparators, that had haemorrhagic events of any severity as outcome.

Two independent investigators conducted the literature search; the revision and the selection of the studies were performed by the working group based on title/abstract and subsequently on full text. Study screening flow diagram is reported in Figure 1.

Since in the Position Paper AISF/SIMI selection criteria were not specified, we assumed they were the same ones we applied in our literature search. Therefore, the 15 studies, which met the inclusion criteria,⁹⁻²³ were analysed together with those identified by the working group AISF/SIMI.²⁴⁻⁶⁸

3 | PROCEDURE-RELATED BLEEDING RISK IN DIFFERENT SETTINGS

Table 1 summarizes the studies included in the analysis.

3.1 | Paracentesis

In clinical practice, paracentesis is usually performed in cirrhotic patients with significant portal hypertension and thrombocytopenia. However, according to the literature, the incidence of post-paracentesis haemorrhagic events was extremely low, and since the presence of portal hypertension is associated with bleeding regardless of platelet count, it was probably related to the patient clinical condition rather than the platelet count.^{10,24-28} Even in the most numerous samples with a clear evaluation of platelet count, no bleeding was recorded in paracentesis performed with platelet count $< 50 \times 10^9/L$.^{10,25}

3.2 | Liver biopsy

Bleeding risk associated with percutaneous liver biopsy was about 0.6% in different studies including numerous sample




FIGURE 1 Study screening flow diagram

sizes, but also very heterogeneous populations in terms of stage of liver disease.^{11,12,32,33} Due to the fact that the presence of severe thrombocytopenia proxies advanced liver disease, thus obviating the need for histological confirmation of the presence of cirrhosis, and to the perceived potential bleeding risk, in clinical practice percutaneous liver biopsy is usually performed in patients without portal hypertension and platelet count $> 50 \times 10^9/L$. The HALT-C trial³³ represented the largest sample of patients with advanced liver disease who underwent percutaneous liver biopsy. Even if the bleeding complications were rare (overall haemorrhagic rate = 0.6%), the study highlighted an increased risk of post-procedural bleeding in patients with platelet count $\leq 60 \times 10^9/L$ (4/76; 5.3%) compared to patients with a platelet count $> 60 \times 10^9/L$ (11/2578; 0.4%), even though in this study a platelet count $< 50 \times 10^9/L$ was an exclusion criterion.

Transjugular liver biopsy is a procedure related to the operator expertise and in clinical practice is usually performed in patients with advanced liver disease, portal hypertension

and thrombocytopenia. In spite of this, bleeding rate from studies was $< 2\%$ and was mainly represented by the occurrence of haematoma at the site of insertion.³⁴⁻³⁷ None of the studies evaluated the association between platelet count and post-procedural bleeding rate.

3.3 | Dentistry

Most of the evidence on this topic is provided by retrospective studies in which bleeding risk seemed to be inherently related to the procedure, or the number of teeth extraction, rather than to platelet count.^{14,38,39} Furthermore, the study of Ward et al³⁸ was severely biased by massive transfusions before the procedure, thus making unfeasible any interpretation of the results in regard to the potential association between bleeding and severe thrombocytopenia. An association between platelet count and post-procedural bleeding was found in the study of Cocero et al,¹³ in which the haemorrhagic rate was 0.4% for patients with platelet count $> 40 \times 10^9/L$ and

TABLE 1 Summary of the studies included in the analysis

Author, year	Study design	Procedures/ patients (n.)	PLT count or PLT cut-off	Findings
Paracentesis				
Webster et al, 1996 ²⁴	Retrospective	179 outpatients	Not specified	4 haemorrhagic complications in patients with $PLT > 80 \times 10^9/L$
Grabau et al, 2004 ²⁵	Retrospective	1100 in 628 pts	$PLT < 50 \times 10^9/L$ in 55.64% of procedures	No bleedings in procedures performed with $PLT < 50 \times 10^9/L$
Pache et al, 2005 ²⁶	Retrospective	4729 paracenteses	Not specified	Severe haemorrhagic complications (6 haemoperitoneum, 3 abdominal wall haematoma) in 0.2% of procedures without association with PLT count
Lin et al, 2005 ²⁷	Prospective observational	410 in 163 pts	$PLT < 50 \times 10^9/L$ in 13% of procedures	Minor bleeding rate (1 local ecchymosis, 1 cutaneous bleeding) in 0.5% of procedures in patients with $PLT = 50-100 \times 10^9/L$
De Gottardi et al, 2009 ²⁸	Prospective observational	515 in 171 pts	$PLT < 50 \times 10^9/L$ in 10% and $PLT < 100 \times 10^9/L$ in 40% of pts	Association between $PLT < 50 \times 10^9/L$ and increased risk of overall complications ($P = .07$). Association with bleeding risk not reported
Rowley et al, 2019 ¹⁰	Retrospective	3116 in 123 pts	$PLT < 50 \times 10^9/L$ in 12% of pts	Overall bleeding rate: 0.2%. No bleeding with $PLT < 50 \times 10^9/L$
Liver biopsy				
Piccinino et al, 1986 ²⁹	Retrospective	68 276 percutaneous biopsies	$PLT > 50 \times 10^9/L$ in all biopsies	Overall rate of major haemorrhagic events: 0.06%. Association between bleeding and PLT not evaluated
Caturelli et al, 1996 ³⁰	Retrospective	NR (only abstract available)	Not specified	Overall rate of haemorrhagic complications: 0.13%. Association between bleeding and PLT not evaluated
Actis et al, 2007 ³¹	Retrospective	835 pts	Not specified	Overall rate of major haemorrhagic events: 0.12%. Association between bleeding and PLT not evaluated
West et al, 2010 ³²	Retrospective	61 187 pts	Not specified	Overall rate of major haemorrhagic events: 0.65%. Association between bleeding and PLT not evaluated
Seeff et al, 2010 ³³	Retrospective	2740 percutaneous biopsies	Pts with $PLT < 50 \times 10^9/L$ were excluded	Overall haemorrhagic rate: 0.6%. Pts with $PLT = 50-60 \times 10^9/L$ was significantly higher than pts with $PLT > 60 \times 10^9/L$ (5.3% vs 0.4%).
Kalambokis et al, 2007 ³⁴	Review	7649 transjugular biopsies in 7189 pts	Cut-off $60 \times 10^9/L$	Haemorrhagic rate $< 2\%$ (minor bleeding). No association with PLT count
Alessandria et al, 2008 ³⁵	Retrospective	306 transjugular biopsies	Not specified	No major complications. No association between bleeding rate and PLT count
Mammen et al, 2008 ³⁶	Retrospective	601 transjugular biopsies	$PLT < 60 \times 10^9/L$ in 20.3% of pts	Haemorrhagic rate = 0.9%. No association with PLT count
Procopet et al, 2012 ³⁷	Prospective	75 transjugular biopsies	Not specified	Haemorrhagic rate = 1.3%. No association with PLT count
Takyar et al, 2017 ¹¹	Retrospective	3357 percutaneous biopsies	Cut-off $100 \times 10^9/L$	Haemorrhagic rate: 0.69% (fatal in 0.09%). $PLT < 100 \times 10^9/L$ was an independent risk factors for post-biopsy bleeding, but % pts with $PLT < 60 \times 10^9/L$ were not different between groups

(Continues)

TABLE 1 (Continued)

Author, year	Study design	Procedures/ patients (n.)	PLT count or PLT cut-off	Findings
Potretzke et al, 2018 ¹²	Retrospective	1876 percutaneous biopsies in 1732 pts	Cut-off $70 \times 10^9/L$	Haemorrhagic rate = 0.69%. No association with PLT count
Dentistry				
Ward et al, 2006 ³⁸	Retrospective	35 procedures in 30 pts	Cut-off $35\text{--}50 \times 10^9/L$ (depending on the risk group)	No association between PLT count and prolonged postoperative bleeding
Perdigao et al, 2012 ³⁹	Prospective observational	35 procedures in 23 pts	$PLT < 50 \times 10^9/L$ in 34% of pts	1 postoperative bleeding (2.9%) in pts with $PLT = 50 \times 10^9/L$. No haemorrhagic complication during procedures (n = 12) with $PLT = 30\text{--}49 \times 10^9/L$
Cocero et al, 2017 ¹³	Retrospective	1,183 extractions in 381 pts	Cut-off $PLT \leq 40 \times 10^9/L$	Haemorrhagic rate: 0.4% in pts with $PLT > 40 \times 10^9/L$ and $INR < 2.5$; 5.88% in pts with $PLT \leq 40 \times 10^9/L$
Medina et al, 2018 ¹⁴	Retrospective	190 extractions	$PLT < 150 \times 10^9/L$ in 96.3% of pts	Overall haemorrhagic rate: 6.3%. Intra-operative bleeding was associated with low count of platelets. However, this counting could explain only 16% of the cases of bleeding.
Endoscopic variceal ligation				
Vieira da Rocha et al, 2009 ⁴⁰	Prospective observational	150 pts	$PLT < 50 \times 10^9/L$ in 12% of pts	Severe post-procedural ulcer bleeding in 7.33% of pts. Risk of bleeding was not associated with PLT count.
Vanbiervliet et al, 2010 ⁴¹	Retrospective	837 ligations in 605 pts	Not specified	Post-procedural bleeding rate: 2.75%. No association between PLT count and bleeding but high platelet ratio index was an independent predictive factor of bleeding
Endoscopic polypectomy				
Jeon et al, 2012 ⁴²	Retrospective	66 in 30 pts	Not specified	Post-procedural bleeding in 3% of procedures. No association between bleeding and PLT count
Lee et al, 2014 ⁴³	Retrospective	89 pts w/ liver cirrhosis + 348 w/o liver disease	Not specified	Post-procedural bleeding in 14.61% of pts. Association between bleeding and PLT not evaluated
Soh et al, 2020 ¹⁵	Retrospective	1267 patients	Cut-off $50 \times 10^9/L$	Haemorrhagic rate (immediate + delayed): 7.5%. $PLT < 50 \times 10^9/L$ significantly associated with immediate post-procedural bleeding (rate: 27.5%; OR = 6.6)
Percutaneous ablation				
Cammà et al, 2005 ⁴⁴	Retrospective	202 pts	$\geq 40 \times 10^9/L$ in all pts	Haemorrhagic rate: 0.50%. Association between bleeding and PLT not evaluated
Livraghi et al, 2008 ⁴⁵	Retrospective	218 pts	$\geq 40 \times 10^9/L$ in all pts	Haemorrhagic rate: 0.92%. Association between bleeding and PLT not evaluated

(Continues)

TABLE 1 (Continued)

Author, year	Study design	Procedures/ patients (n.)	PLT count or PLT cut-off	Findings
Goto et al, 2010 ⁴⁶	Retrospective	4133 in 2154 pts	Mean PLT count = $125 \pm 33 \times 10^9/L$ (50-669)	Haemorrhagic complications rate: 1.5%. Low PLT count was a significant risk factor for haemoperitoneum (PLT $\geq 50 \times 10^9/L$ was an inclusion criteria)
Park et al, 2017 ¹⁶	Retrospective	1843 in 1211 patients	Mean PLT count = $140 \pm 85 \times 10^9/L$	Post-procedural bleeding rate was 0.6%, and the risk was significantly greater in patients with PLT $< 50 \times 10^9/L$ (OR = 8.79)
Liver transplantation				
McCluskey et al, 2006 ⁴⁷	Retrospective	460 pts	Not specified	Incidence of massive blood transfusion: 42%. PLT $< 70 \times 10^9/L$ was an independent predictor of massive blood transfusion (+32% vs PLT $> 70 \times 10^9/L$)
Massicotte et al, 2008 ⁴⁸	Prospective + retrospective	200 pts	$< 50 \times 10^9/L$ in 18% of pts; $< 30 \times 10^9/L$ in 4% of pts	No association between PLT count and transfusion rate
Massicotte et al, 2012 ⁴⁹	Retrospective	503 pts	Not specified	No significant association between PLT count and blood loss
Esmat Gamil et al, 2012 ⁵⁰	Prospective observational	286 pts	Not specified	No significant association between PLT count and blood loss
Li et al, 2014 ¹⁷	Retrospective	241 pts	Not specified	Postoperative bleeding in 4.98% of pts. No significant association between PLT count and bleeding risk
Akamatsu et al, 2015 ¹⁸	Retrospective	403 pts	Mean PLT count $86 \pm 70 \times 10^9/L$	Haemorrhagic episodes in 8.68% of pts. No significant association between PLT count and blood loss
Eghbal et al, 2019 ¹⁹	Retrospective	754 pts	Not specified	PLT count was inversely correlated with total bleeding
Liver surgery				
Wei et al, 2003 ⁵¹	Retrospective	155 pts	Median PLT count $205 \times 10^9/L$ (82-473)	Postoperative intra-abdominal haemorrhage in 5% of patients. Association between bleeding and PLT not evaluated
Kubo et al, 2007 ⁵²	Retrospective	100 pts	Not specified	Postoperative bleeding in 4% of patients. Association between bleeding and PLT not evaluated
Palavecino et al, 2009 ⁵³	Retrospective	1557 resections in 1477 pts	Median PLT count $232 \times 10^9/L$ (64.0-775.0)	PLT $< 100 \times 10^9/L$ (1% of pts) was an independent risk factors for blood transfusion (OR = 8.8)
Hsu et al, 2009 ⁵⁴	Retrospective	1027 resections	Not specified	PLT $< 100 \times 10^9/L$ was correlated with perioperative mortality in the univariate analysis but not in the multivariate one
Cockbain et al, 2010 ⁵⁵	Retrospective	589 pts	Cut-off $150 \times 10^9/L$	No association between PLT $> 150 \times 10^9/L$ and lower transfusion rate
Yang et al, 2011 ⁵⁶	Retrospective	305	Not specified	Haemorrhagic complications in 2.62% of pts. PLT count $< 100 \times 10^9/L$ was independently correlated with postoperative morbidity and hospital mortality

(Continues)

TABLE 1 (Continued)

Author, year	Study design	Procedures/ patients (n.)	PLT count or PLT cut-off	Findings
Vascular catheter insertion				
Fisher et al, 1999 ⁵⁷	Retrospective	658 cannulations in 283 pts	PLT < $50 \times 10^9/L$ in ~ 25% pts	1 haemothorax in pts with PLT = $68 \times 10^9/L$. PLT $\leq 10 \times 10^9/L$ significantly associated with superficial haematoma vs PLT > $50 \times 10^9/L$ (4,8% vs. 1,6%, respectively)
Estcourt et al, 2015 ²⁰	Systematic review	-	Not specified	No evidence from RCTs to determine whether PLT transfusions are required prior to central line insertion in patients with thrombocytopenia, and, if a PLT transfusion is required, what is the correct threshold.
HVPG measurement				
Bosch et al 2009 ⁵⁸	Review + single-centre experience	12 000 measurement	Not reported	Major complications have been limited to local injury at the puncture site and include leakage, haematoma and rarely fistulae or Horner syndrome
Woolfson et al, 2013 ⁵⁹	Retrospective	52 HVPG measurements in 49 children	PLT < $100 \times 10^9/L$ in 28 pts	Variceal bleeding and variceal bleeding + ascites occurred each in 1/7 patients with cirrhosis. Association between bleeding and PLT not evaluated
Cholecystectomy				
Sleeman et al, 1998 ⁶⁰	Retrospective	25 pts	PLT < $100 \times 10^9/L$ in 36%	Association between bleeding and PLT not evaluated
Da Silveira et al 2006 ⁶¹	Retrospective	99 pts	Not reported	Association between bleeding and PLT not evaluated
Delis et al, 2010 ⁶²	Retrospective	220 procedures	Transfusion when PLT < $50 \times 10^9/L$	Intra-operative bleeding rate: 8%. Association between bleeding and PLT not evaluated
Herniotomy				
Carbonell et al, 2005 ⁶³	Retrospective	32 033 procedures	Not reported	Association between bleeding and PLT not evaluated.
Ammar et al, 2010 ⁶⁴	Prospective	80 pts	Not reported	Association between bleeding and PLT not evaluated
Thoracentesis				
Castellote et al, 2001 ⁶⁵	Retrospective	245 thoracentesis in 69 cirrhotic pts	Not reported	Haemorrhagic rate: 2%. Association between bleeding and PLT not evaluated
Xiol et al, 2001 ⁶⁶	Retrospective	215 thoracenteses in 60 cirrhotic pts	Not reported	Association between bleeding and PLT not evaluated
Urological surgery				
Nielsen et al, 2001 ⁶⁷	Retrospective	180 pts	Not reported	Association between bleeding and PLT not evaluated
Lund et al, 2003 ⁶⁸	Retrospective	611	Not reported	Association between bleeding and PLT not evaluated

(Continues)

TABLE 1 (Continued)

Author, year	Study design	Procedures/ patients (n.)	PLT count or PLT cut-off	Findings
Miscellaneous				
Shah et al, 2015 ²¹	Prospective observational	380 pts	Cut-off $50 \times 10^9/L$	Clinically significant bleeding following high-risk procedures occurred in 3 patients with significant coagulopathy and 0 patients without significant coagulopathy ($P = .061$)
De Pietri et al, 2016 ⁹	RCT open-label ITT	60 pts	Cut-off $50 \times 10^9/L$	Bleeding occurred in 1.7% of patients (1/60) following paracentesis (PLT = $111 \times 10^9/L$)
Napolitano et al, 2017 ²²	Prospective observational	852 procedures in 363 pts	Cut-off $50 \times 10^9/L$	Overall bleeding complication rate: 2.75%. No bleeding in 90 procedures with PLT count $< 50 \times 10^9/L$ and no association were identified between PLT count and bleeding risk
Vuyyuru et al, 2020 ²³	RCT open-label	60 pts	Cut-off $50 \times 10^9/L$	No bleeding in 58 procedures with PLT count $< 50 \times 10^9/L$

Abbreviations: HVPg, hepatic venous pressure gradient; INR, international normalized ratio; OR, odds ratio; PLT, platelet; pts, patients.

international normalized ratio (INR) < 2.5 , and 5.88% in patients with platelet count $\leq 40 \times 10^9/L$. Finally, in the only prospective study³⁹ postoperative bleeding occurred in only one procedure (2.9%) performed in a patient with liver cirrhosis, INR = 2.50 and platelet count = $50 \times 10^9/L$, whereas no bleeding occurred in procedures performed in patients with platelets = $30\text{--}49 \times 10^9/L$.

3.4 | Endoscopic variceal ligation

In the two studies analysed, the post-procedural bleeding rate ranged from 2.75% in the case-control study of Vanbiervliet et al,⁴¹ to 7.33% in the prospective study of Viera da Rocha et al⁴⁰. In both cases, there was no association between bleeding risk and platelet count. In general, post-ligation bleeding was related to technical problem occurred during the procedure, late bleeding or portal hypertension.

3.5 | Endoscopic polypectomy

All the studies identified were retrospective and potentially biased by the heterogeneity of the investigated population including both cirrhotic and noncirrhotic patients.^{15,42,43} Only the study by Soh et al¹⁵ identified a correlation between post-procedural bleeding and platelet count: while the overall haemorrhagic rate was 7.5%, in patients with platelets $< 50 \times 10^9/L$, the immediate post-procedural bleeding rate was 27.5% with a relative risk of about 6.

3.6 | Percutaneous radio-frequency ablation of hepatocellular carcinoma

In clinical practice, percutaneous radio-frequency ablation of hepatocellular carcinoma (HCC) is rarely performed in patients with platelets $< 50 \times 10^9/L$ and is usually preceded by platelet transfusions and a close monitoring of platelet count. Therefore, the rate of bleeding following the radio-frequency ablation of HCC was lower than 1%.^{16,44,45} Only the study of Park et al¹⁶ found a correlation between a platelet count $< 50 \times 10^9/L$ and an increased risk of post-procedural bleeding rate (OR = 8.79). However, the study was biased by prophylactic platelet transfusion in patients with platelets $< 50 \times 10^9/L$. Finally, the study of Goto et al⁴⁶ showed a haemorrhagic complication rate of 1.5% in 4133 radio-frequency ablations (not only HCC), and thrombocytopenia was identified as significant risk factor for hemoperitoneum, even though patients with severe thrombocytopenia (platelets $< 50 \times 10^9/L$) were not included in the study.

3.7 | Liver transplantation

The risk and extent of bleeding during liver transplantation were difficult to quantify and were generally reported only as indirect evidence (i.e. number of transfused blood products or amount of blood loss). None of the studies showed an association between platelet count and intra- or post-transplantation bleeding.^{17-19,47-50} Indeed, in this setting, the bleeding risk cannot be evaluated on the basis of blood

coagulation parameters, since it may be influenced by other recipient's conditions, technical difficulties and portal hypertension control. Besides improvements in surgical experience and techniques in liver transplantation, strategies to reduce the use of blood products termed 'patient blood management' are increasingly adopted. For monitoring of haemostasis disturbances, thromboelastography (TEG) or thromboelastometry (TEM) is indicated as the best blood tests that can guide the application of plasma components, platelets and antifibrinolytics.⁶⁹

3.8 | Liver surgery

In liver surgery, portal hypertension is the main determinant of outcome; in a large series published in 2011, even mild thrombocytopenia (platelet count of less than $150 \times 10^9/L$) predicted major postoperative complications and mortality after resection of HCC independently of functional scores such as Child-Pugh or MELD score.⁷⁰

However, in this setting, bleeding rate and risk factors were very difficult to identify due to heterogeneous populations (i.e. inclusion of cirrhotic and noncirrhotic patients). All the studies were retrospective, and none evaluated the association between platelet count and bleeding risk in liver surgery.⁵¹⁻⁵⁶ This was probably due to the fact that in clinical practice moderate-to-severe thrombocytopenia is often considered a contraindication to liver surgery and patients are treated with pre- or intra-operative platelet transfusions.

3.9 | Abdominal surgery and other invasive procedures

The vascular catheter insertion and the hepatic venous pressure gradient measurement are procedures related to the operator expertise and are usually performed in patients at high risk of bleeding due to advanced liver disease, portal hypertension and thrombocytopenia. However, the available studies were not sufficient to determine a relationship between platelet count and bleeding risk following these two procedures.^{20,57-59}

Regarding cholecystectomy and herniotomy, the wide heterogeneity in the management of blood coagulation parameters in the pre-procedural phases made the relationship between thrombocytopenia and haemorrhagic risk not evaluable. Furthermore, the available studies did not evaluate this association.⁶⁰⁻⁶⁴ Finally, also the available evidence related to thoracentesis^{65,66} and urological surgery^{67,68} was not sufficient to assess the association between platelet count and post-procedural bleeding risk.

3.10 | Miscellaneous

Some studies evaluated the overall risk of bleeding in cirrhotic patients submitted to different procedures and the association with platelet count and/or coagulopathy.^{9,13,15,23} In the open-label, intention-to-treat trial of De Pietri et al⁹ cirrhosis and significant coagulopathy (defined as $INR > 1.8$ and/or platelet count $< 50 \times 10^9/L$) did not expose to an increased procedure-related bleeding risk, regardless of the procedure (i.e. high- or low-risk procedures), although the cohort included was small (i.e. 30 patients per arm) and all patients with severe thrombocytopenia in the standard of care arm received prophylactic platelet transfusions.

Similarly, the prospective case series of Napolitano et al¹⁵ did not identify any association between platelet count and post-procedural bleeding risk, not even in patients with a platelet count $< 50 \times 10^9/L$. In this case, the only parameter associated with the risk of bleeding was the number of invasive procedures sequentially performed in each single patient: 3 events following 598 single procedures (bleeding rate: 0.5%) and 7 events following multiple procedures (1.5%).¹⁵

Also, in the randomized controlled trial of Vuyyuru et al²³ no bleeding complications occurred following 58 procedures in cirrhotic patients with platelet count $< 50 \times 10^9/L$, whereas in the prospective multicentre study of Shah et al,¹³ in which none of the patients received peri-procedural correction of abnormal coagulation parameters, the occurrence of clinically significant bleeding following high-risk procedures (i.e. cholecystectomy, splenectomy, chemoembolization, central vein cannulation, percutaneous liver biopsy and endoscopic polypectomy) tended to be greater in patients with significant coagulopathy (defined as $INR \geq 1.5$ and/or platelet count $\leq 50 \times 10^9/L$) as compared to patients without significant coagulopathy (3 vs 0, $P = .061$), although it was not possible to single out the role played by thrombocytopenia alone on the bleeding risk in this study.

4 | INTERPRETATION OF THE RESULTS AND POTENTIAL CLINICAL IMPLICATIONS

Despite the lack of solid evidence identified in the previous consensus conference and the call for prospective studies to address the issue of procedure-related bleeding risk in patients with liver disease, even most of the new studies included in our analysis had the limitation of not adequately assessing the association between post-procedural bleeding risk and platelet count in patients with chronic liver disease. We also found other limitations of the available literature. The first is that the majority of studies that investigated the role of platelet count were retrospective and heterogeneous in terms

of population (i.e. inclusion of cirrhotic and noncirrhotic patients). Also, it is important to note that in clinical practice moderate-to-severe thrombocytopenia is often considered a contraindication to some procedures (e.g. percutaneous radio-frequency ablation of HCC, liver biopsy and liver surgery) and that patients are frequently treated with plasma and pre- or intra-operative platelet transfusion in order to mitigate the risk of bleeding.^{8,71-74} Therefore, it could be possible that some results were biased by these prophylactic interventions.

Only a few studies, among those who assessed the risk of bleeding in relation to platelet count, found that thrombocytopenia may be predictive of bleeding following percutaneous liver biopsy,^{11,33} dental extractions,^{13,14} percutaneous ablation of liver tumours^{16,46} and endoscopic polypectomy.¹⁵ Noteworthy, none of the prospective studies included in this review highlighted a significant correlation between post-procedural haemorrhagic rate and platelet count.^{9,21-23}

Despite the above limitations, that would require the conduction of prospective studies properly designed to evaluate the bleeding risk in patients with chronic liver disease undergoing invasive procedures, according to platelet count, the available literature highlighted that severe thrombocytopenia is one of the most frequent issues to exclude cirrhotic patients to invasive procedures, which could be, in some cases, life-saving, such as percutaneous radio-frequency ablation in malignant lesions. However, available evidence had also the strength to confirm that there is no platelet count threshold at which bleeding is predictable, as other factors contribute to bleeding risk.

Indeed, it has been shown that in patients with chronic liver disease, even at an advanced stage, platelet count alone cannot be considered the only predictor of increased risk of bleeding,^{75,76} while platelet count should be properly considered in the presence of other risk factors such as sepsis and acute kidney injury, in order to provide a more accurate estimate of the bleeding risk of the patients.^{21,77}

In addition, in cirrhotic patients the aetiology and severity of the disease can influence the haemostatic balance and comorbidities could alter the feeble haemostatic equilibrium in patients with advanced liver disease.^{8,77}

Despite the limited evidence available, several Position Papers and Guidelines of Scientific Societies recommend the correction of thrombocytopenia in patients with chronic liver disease and platelet count $< 50 \times 10^9/L$ who are scheduled to undergo invasive procedures.^{8,71-74} In these patients, although there is no solid evidence of its efficacy on raising and maintaining an adequate platelet count level,^{22,78} the standard of care is platelet transfusion. On the contrary, in patients with advanced liver disease there is evidence that prophylactic blood products transfusions, and the resultant volume expansion in a short timeframe, may aggravate portal hypertension and therefore paradoxically determine an increase in bleeding risk.⁷⁹ In this context, in cirrhotic patients with high risk of bleeding, thrombopoietin receptor agonists (TPO-RAs) may represent an advantageous therapeutic alternative.⁸⁰⁻⁸² TPO-RAs may improve patient clinical management as they are able to increase patient's platelet counts in a predictable fashion, thus allowing to plan invasive procedures and avoiding





FIGURE 2 Invasive procedures performed in the lusutrombopag group during the (A) L-PLUS 1 and (B) L-PLUS 2 studies. APC = argon plasma coagulation; EIS = endoscopic injection sclerosis; EVL = endoscopic variceal ligation; GI = gastrointestinal; MCT = microwave coagulation therapy; PEIT = percutaneous ethanol injection therapy; RFA = radio-frequency ablation; TACE = transcatheter arterial chemoembolization. (A) From: Hidaka H et al. Lusutrombopag Reduces Need for Platelet Transfusion in Patients With Thrombocytopenia Undergoing Invasive Procedures. Clin Gastroenterol Hepatol. 2019;17(6):1192-1200 (B) Elaborated from: Peck-Radosavljevic et al Lusutrombopag for the Treatment of Thrombocytopenia in Patients With Chronic Liver Disease Undergoing Invasive Procedures (L-PLUS 2). Hepatology. 2019;70(4):1336-1348

FIGURE 3 Median PCs over time for patients treated with lusutrombopag (without platelet transfusion) or placebo (with platelet transfusion) in the (A) PLUS-1 and (B) PLUS-2 studies. From: (A) Hidaka H et al's Lusutrombopag Reduces Need for Platelet Transfusion in Patients With Thrombocytopenia Undergoing Invasive Procedures. *Clin Gastroenterol Hepatol*. 2019;17(6):1192-1200 (B) Peck-Radosavljevic et al Lusutrombopag for the Treatment of Thrombocytopenia in Patients With Chronic Liver Disease Undergoing Invasive Procedures (L-PLUS 2). *Hepatology*. 2019;70(4):1336-1348



the risk of postponement or cancellation of procedures due to an inadequate increase in platelet count, as is often observed with platelet transfusions.⁸⁰⁻⁸² However, the use of TPO-RAs has been associated with venous thromboembolism in patients with chronic liver disease, probably because of too high a rise in the platelet count and platelet hyperactivity in liver cirrhosis patients.⁷⁵ More in detail, as reported by Loffredo et al,⁸³ a statistically significant association between thrombotic risk and TPO-RAs use was observed only in patients treated with eltrombopag.

Therefore, after the early termination of the clinical trial of eltrombopag, due to the increased thrombotic risk,⁸⁴ lusutrombopag was the first oral drug approved by EMA for the treatment of severe thrombocytopenia in patients with chronic liver disease undergoing invasive procedures.⁸⁵ Lusutrombopag showed efficacy in reducing the need for platelet transfusion, raising the platelet count $> 50 \times 10^9/L$ at the time of procedures and maintaining an adequate platelet level following the procedures, thus granting a safe lingering effect that may theoretically protect from delayed bleeding or allow to perform repeated invasive procedures.^{80,81} In the pivotal studies L-PLUS 1 and L-PLUS 2, in fact, lusutrombopag

allowed to reach the platelet threshold of $50 \times 10^9/L$ in about 9 days, to maintain it for a median of 20.9 days, and safely performed invasive procedures (Figures 2 and 3).^{80,81}

Furthermore, the aggregated data showed that the use of lusutrombopag was associated with a numerical lower rate of post-procedural bleeding (6.7% vs 10.6%) without increased risk of thrombosis.^{80,81,85} Platelet increase with lusutrombopag was in fact more moderate than with other TPO-RAs (median highest platelet count eltrombopag vs lusutrombopag: $140 \times 10^9/L$ vs $80 \times 10^9/L$).^{80,84}

In the ADAPT-1 and ADAPT-2, pivotal studies on avatrombopag patients with low ($< 40 \times 10^9/L$) and high ($40-50 \times 10^9/L$) baseline platelet count received avatrombopag 40 and 60 mg, respectively.⁸² In both cohorts, the proportion of patients who did not require a platelet transfusion after randomization and up to 7 days after the procedure was higher in those who received avatrombopag, compared with placebo. Moreover, in both avatrombopag treatment groups, platelet count increase was observed from day 4, reaching a maximum at days 10-13. The mean platelet count remained at or above $50 \times 10^9/L$ at day 17, and only 3 patients reach platelet count $> 200 \times 10^9/L$.⁸²

5 | CONCLUSIONS AND FUTURE DIRECTIONS

Despite several studies were conducted in the past, there is still a lack of adequate and solid data depicting the risk of bleeding following invasive procedures in patients with advanced liver disease, and its potential association with decreased platelet count. This notwithstanding, the best evidence currently available points to an association between severe thrombocytopenia and an increased risk of bleeding in patients with advanced cirrhosis undergoing procedures, in particular in subjects who undergo 'closed procedures' such as biopsies of parenchymal organs or liver tumour ablations. In this regard, international guidelines suggest that severe thrombocytopenia should be corrected before procedures in these patients, nevertheless, due to the lack of literature to support definitive conclusions about the appropriate target platelet counts to improve the risk of bleeding, there is an enormous variability in the use of prophylactic platelet transfusions.⁸⁶ It is, however, well established that the prophylactic use of platelet transfusions for these patients is of unpredictable efficacy and biased by potential adverse events including transfusion reactions, sepsis, refractoriness to further platelet transfusions, prolonged hospitalization and increased costs. Specifically, refractoriness to further platelet transfusions may add complexity to the management of these patients and reduce treatment options for bleeding associated with invasive procedures and/or surgery including liver transplant.

For all the above, research for therapeutic options alternative to platelet transfusion is welcome and TPO-RAs seem to represent a valid option given the safety and efficacy, simplifying the clinical management of these patients. It is important to underline that, unlike platelet transfusion, the use of TPO-RAs represents the only strategy capable of obtaining a real significant increase in the platelet count. Furthermore, the use of TPO-RAs may be associated with the improvement in global healthcare resource utilisation, as blood product transfusions—and in particular platelets—are quite often used in clinical practice to increase platelets in patients undergoing procedures, and therefore, in this setting a treatment alternative may increase platelet availability for other clinical purposes.^{86,87}

In this context, still to be detailed is for which invasive procedure TPO-RA prescription can be allocated and if some liver-related disease complication may question the therapeutic efficacy. Despite a clear indication of the use of platelet growth factor in cirrhotic patients undergoing procedures at particular risk of bleeding, a valid therapeutic address to be considered is in liver transplant candidates, although in advanced stage of Child-Pugh score the safety of TPO-RAs has not yet been assessed. Routine procedures such as dental extraction, endoscopic polypectomy,

ligation of oesophageal varices, transjugular intrahepatic portosystemic shunt (TIPS) and HCC transarterial chemoembolization (TACE) could benefit from a single period of drug therapy with the maximum result in terms of lower flow overload (compared with platelet transfusion), fewer days of hospitalization (due to ineffective increase platelet with transfusions) and lower risk of bleeding due to the achievement of a more adequate level of platelets in the plasma by using platelet growth factors.

In the next future, well-designed studies may disclose whether the use of TPO-RAs may actually be associated with a decreased risk of bleeding following procedures in patients with liver disease as compared to platelet transfusions, although planning such studies may represent a difficult task due to the variability of clinical situations, the vast array of potential procedures and their different risk of bleeding, and the overall low risk of bleeding that should call for the enrolment of very large cohorts of patients. Furthermore, real-life data could add important information on the effectiveness and safety of TPO-RAs in the management of invasive procedures in cirrhotic patients at high risk of bleeding, thus providing the basis of a potential new standard of care.

ACKNOWLEDGEMENTS

We would like to thank Ombretta Bandi, from SEEd Medical Publishers, who provided writing assistance and journal styling services, and Lorenzo Pradelli, from AdRes—Health Economics & Outcomes Research, who performed literature search. Editorial services and literature search were funded by Shionogi srl. Shionogi srl had no role in study design, nor in collection, analysis and interpretation of data.

CONFLICT OF INTEREST

DA declares research grant and consultant fee from Intercept Pharma, research grant from Vesta (USA) and consultant fee from Aboca. NC declares consultant fee from Shionogi. EGG declares consultant fee from Shionogi, BMS, Gilead Sciences, AbbVie, Bayer and GSK. AI and PT have nothing to disclose. MM declares consultant fee from Shionogi, Gilead Sciences, AbbVie and Biotest. FV declares consultant fee from Shionogi.

AUTHOR CONTRIBUTIONS

The review topic was proposed by all authors. Literature selection and analysis of data were undertaken by all authors. Critical revisions of the manuscript were conducted by all authors. Final approval for the version to be published was obtained from all authors. All authors read and approved the final manuscript.

ORCID

Edoardo Giovanni Giannini  <https://orcid.org/0000-0001-8526-837X>

REFERENCES

- Qamar AA, Grace ND, Groszmann RJ, et al. Incidence, prevalence, and clinical significance of abnormal hematologic indices in compensated cirrhosis. *Clin Gastroenterol Hepatol*. 2009;7(6):689-695.
- Louie KS, Micallef JM, Pimenta JM, Forssen UM. Prevalence of thrombocytopenia among patients with chronic hepatitis C: a systematic review. *J Viral Hepat*. 2011;18(1):1-7.
- Peck-Radosavljevic M. Thrombocytopenia in chronic liver disease. *Liver Int*. 2017;37(6):778-793.
- Giannini EG. Review article: thrombocytopenia in chronic liver disease and pharmacologic treatment options. *Aliment Pharmacol Ther*. 2006;23(8):1055-1065.
- Bashour FN, Teran JC, Mullen KD. Prevalence of peripheral blood cytopenias (hypersplenism) in patients with nonalcoholic chronic liver disease. *Am J Gastroenterol*. 2000;95(10):2936-2939.
- Afdhal N, McHutchison J, Brown R, et al. Thrombocytopenia associated with chronic liver disease. *J Hepatol*. 2008;48(6):1000-1007.
- Giannini EG, Greco A, Marengo S, Andorno E, Valente U, Savarino V. Incidence of bleeding following invasive procedures in patients with thrombocytopenia and advanced liver disease. *Clin Gastroenterol Hepatol*. 2010;8(10):899-e109.
- Andriulli A, Tripodi A, Angeli P, et al. Under the auspices of the Italian Association for the Study of Liver Diseases (AISF) and the Italian Society of Internal Medicine (SIMI). Hemostatic balance in patients with liver cirrhosis: report of a consensus conference. *Dig Liver Dis*. 2016;48(5):455-467.
- De Pietri L, Bianchini M, Montalti R, et al. Thrombelastography-guided blood product use before invasive procedures in cirrhosis with severe coagulopathy: a randomized, controlled trial. *Hepatology*. 2016;63(2):566-573.
- Rowley MW, Agarwal S, Seetharam AB, Hirsch KS. Real-time ultrasound-guided paracentesis by radiologists: near zero risk of hemorrhage without correction of coagulopathy. *J Vasc Interv Radiol*. 2019;30(2):259-264.
- Takyar V, Etzion O, Heller T, et al. Complications of percutaneous liver biopsy with Klatzkin needles: a 36-year single-centre experience. *Aliment Pharmacol Ther*. 2017;45(5):744-753.
- Potretzke TA, Saling LJ, Middleton WD, Robinson KA. Bleeding complications after percutaneous liver biopsy: do subcapsular lesions pose a higher risk? *Am J Roentgenol*. 2018;211(1):204-210.
- Cocero N, Bezzi M, Martini S, Carossa S. Oral surgical treatment of patients with chronic liver disease: assessments of bleeding and its relationship with thrombocytopenia and blood coagulation parameters. *J Oral Maxillofac Surg*. 2017;75(1):28-34.
- Medina JB, Andrade NS, de Paula EF, et al. Bleeding during and after dental extractions in patients with liver cirrhosis. *Int J Oral Maxillofac Surg*. 2018;47(12):1543-1549.
- Soh H, Chun J, Hong SW, et al. Child-Pugh B or C cirrhosis increases the risk for bleeding following colonoscopic polypectomy [published online ahead of print, 2019 Dec 11]. *Gut Liv*. 2020;14(6):755-764. <https://doi.org/10.5009/gnl19131>
- Park JG, Park SY, Tak WY, et al. Early complications after percutaneous radiofrequency ablation for hepatocellular carcinoma: an analysis of 1,843 ablations in 1,211 patients in a single centre: experience over 10 years. *Clin Radiol*. 2017;72(8):692.e9-692.e15.
- Li C, Wen TF, Yan LN, Li B. Risk factors for abdominal bleeding after living-donor liver transplant. *Exp Clin Transplant*. 2014;12(5):424-428.
- Akamatsu N, Sugawara Y, Nakazawa A, et al. Hemostatic status in liver transplantation: association between preoperative procoagulants/anticoagulants and postoperative hemorrhaging/thrombosis. *Liver Transpl*. 2015;21(2):258-265.
- Eghbal MH, Samadi K, Khosravi MB, et al. The impact of preoperative variables on intraoperative blood loss and transfusion requirements during orthotopic liver transplant. *Exp Clin Transplant*. 2019;17(4):507-512.
- Estcourt LJ, Desborough M, Hopewell S, Trivella M, Doree C, Stanworth S. Comparison of different platelet transfusion thresholds prior to insertion of central lines in patients with thrombocytopenia. *Cochrane Database Syst Rev*. 2015;2015(6):CD011771.
- Shah A, Amarapurkar D, Dharod M, et al. Coagulopathy in cirrhosis: a prospective study to correlate conventional tests of coagulation and bleeding following invasive procedures in cirrhotics. *Indian J Gastroenterol*. 2015;34(5):359-364.
- Napolitano G, Iacobellis A, Merla A, et al. Bleeding after invasive procedures is rare and unpredicted by platelet counts in cirrhotic patients with thrombocytopenia. *Eur J Intern Med*. 2017;38:79-82.
- Vuyyuru SK, Singh AD, Gamanagatti SR, Rout G, Gunjan D, Shalimar. A randomized control trial of thromboelastography-guided transfusion in cirrhosis for high-risk invasive liver-related procedures. *Dig Dis Sci*. 2020;65(7):2104-2111.
- Webster ST, Brown KL, Lucey MR, Nostrant TT. Hemorrhagic complications of large volume abdominal paracentesis. *Am J Gastroenterol*. 1996;91(2):366-368.
- Grabau CM, Crago SF, Hoff LK, et al. Performance standards for therapeutic abdominal paracentesis. *Hepatology*. 2004;40(2):484-488.
- Pache I, Bilodeau M. Severe haemorrhage following abdominal paracentesis for ascites in patients with liver disease. *Aliment Pharmacol Ther*. 2005;21(5):525-529.
- Lin CH, Shih FY, Ma MH, Chiang WC, Yang CW, Ko PC. Should bleeding tendency deter abdominal paracentesis? *Dig Liver Dis*. 2005;37(12):946-951.
- De Gottardi A, Thévenot T, Spahr L, et al. Risk of complications after abdominal paracentesis in cirrhotic patients: a prospective study. *Clin Gastroenterol Hepatol*. 2009;7(8):906-909.
- Piccinino F, Sagnelli E, Pasquale G, Giusti G. Complications following percutaneous liver biopsy. A multicentre retrospective study on 68,276 biopsies. *J Hepatol*. 1986;2(2):165-173.
- Caturelli E, Giacobbe A, Facciorusso D, et al. Percutaneous biopsy in diffuse liver disease: increasing diagnostic yield and decreasing complication rate by routine ultrasound assessment of puncture site. *Am J Gastroenterol*. 1996;91(7):1318-1321.
- Actis GC, Olivero A, Lagget M, Pellicano R, Smedile A, Rizzetto M. The practice of percutaneous liver biopsy in a gastrohepatology day hospital: a retrospective study on 835 biopsies. *Dig Dis Sci*. 2007;52(10):2576-2579.
- West J, Card TR. Reduced mortality rates following elective percutaneous liver biopsies. *Gastroenterology*. 2010;139(4):1230-1237.
- Seeff LB, Everson GT, Morgan TR, et al. Complication rate of percutaneous liver biopsies among persons with advanced chronic liver disease in the HALT-C trial. *Clin Gastroenterol Hepatol*. 2010;8(10):877-883.
- Kalambokis G, Manousou P, Vibhakorn S, et al. Transjugular liver biopsy—indications, adequacy, quality of specimens, and complications—a systematic review. *J Hepatol*. 2007;47(2):284-294.

35. Alessandria C, Debernardi-Venon W, Rizzetto M, Marzano A. Transjugular liver biopsy: a relatively simple procedure with an indefinite past and an expected brilliant future. *J Hepatol*. 2008;48(1):171-173.
36. Mammen T, Keshava SN, Eapen CE, et al. Transjugular liver biopsy: a retrospective analysis of 601 cases. *J Vasc Interv Radiol*. 2008;19(3):351-358.
37. Procopet B, Bureau C, Métivier S, et al. Tolerance of liver biopsy in a tertiary care center: comparison of the percutaneous and the transvenous route in 143 prospectively followed patients. *Eur J Gastroenterol Hepatol*. 2012;24(10):1209-1213.
38. Ward BB, Weideman EM. Long-term postoperative bleeding after dentoalveolar surgery in the pretransplant liver failure patient. *J Oral Maxillofac Surg*. 2006;64(10):1469-1474.
39. Perdigão JP, de Almeida PC, Rocha TD, et al. Postoperative bleeding after dental extraction in liver pretransplant patients. *J Oral Maxillofac Surg*. 2012;70(3):e177-e184.
40. Vieira da Rocha EC, D'Amico EA, Caldwell SH, et al. A prospective study of conventional and expanded coagulation indices in predicting ulcer bleeding after variceal band ligation. *Clin Gastroenterol Hepatol*. 2009;7(9):988-993.
41. Vanbiervliet G, Giudicelli-Bornard S, Piche T, et al. Predictive factors of bleeding related to post-banding ulcer following endoscopic variceal ligation in cirrhotic patients: a case-control study. *Aliment Pharmacol Ther*. 2010;32(2):225-232.
42. Jeon JW, Shin HP, Lee JI, et al. The risk of postpolypectomy bleeding during colonoscopy in patients with early liver cirrhosis. *Surg Endosc*. 2012;26(11):3258-3263.
43. Lee S, Park SJ, Cheon JH, et al. Child-Pugh score is an independent risk factor for immediate bleeding after colonoscopic polypectomy in liver cirrhosis. *Yonsei Med J*. 2014;55(5):1281-1288.
44. Cammà C, Di Marco V, Orlando A, et al. Treatment of hepatocellular carcinoma in compensated cirrhosis with radio-frequency thermal ablation (RFTA): a prospective study. *J Hepatol*. 2005;42(4):535-540.
45. Livraghi T, Meloni F, Di Stasi M, et al. Sustained complete response and complications rates after radiofrequency ablation of very early hepatocellular carcinoma in cirrhosis: is resection still the treatment of choice? *Hepatology*. 2008;47(1):82-89.
46. Goto E, Tateishi R, Shiina S, et al. Hemorrhagic complications of percutaneous radiofrequency ablation for liver tumors. *J Clin Gastroenterol*. 2010;44(5):374-380.
47. McCluskey SA, Karkouti K, Wijesundera DN, et al. Derivation of a risk index for the prediction of massive blood transfusion in liver transplantation. *Liver Transpl*. 2006;12(11):1584-1593.
48. Massicotte L, Beaulieu D, Thibeault L, et al. Coagulation defects do not predict blood product requirements during liver transplantation. *Transplantation*. 2008;85(7):956-962.
49. Massicotte L, Denault AY, Beaulieu D, et al. Transfusion rate for 500 consecutive liver transplantations: experience of one liver transplantation center. *Transplantation*. 2012;93(12):1276-1281.
50. Esmat Gamil M, Pirenne J, Van Malenstein H, et al. Risk factors for bleeding and clinical implications in patients undergoing liver transplantation. *Transplant Proc*. 2012;44(9):2857-2860.
51. Wei AC, Tung-Ping Poon R, Fan ST, Wong J. Risk factors for perioperative morbidity and mortality after extended hepatectomy for hepatocellular carcinoma. *Br J Surg*. 2003;90(1):33-41. <https://doi.org/10.1002/bjs.4018>
52. Kubo S, Takemura S, Yamamoto S, et al. Risk factors for massive blood loss during liver resection for hepatocellular carcinoma in patients with cirrhosis. *Hepatogastroenterology*. 2007;54(75):830-833.
53. Palavecino M, Kishi Y, Chun YS, et al. Two-surgeon technique of parenchymal transection contributes to reduced transfusion rate in patients undergoing major hepatectomy: analysis of 1,557 consecutive liver resections. *Surgery*. 2010;147(1):40-48.
54. Hsu KY, Chau GY, Lui WY, Tsay SH, King KL, Wu CW. Predicting morbidity and mortality after hepatic resection in patients with hepatocellular carcinoma: the role of Model for End-Stage Liver Disease score. *World J Surg*. 2009;33(11):2412-2419.
55. Cockbain AJ, Masudi T, Lodge JP, Toogood GJ, Prasad KR. Predictors of blood transfusion requirement in elective liver resection. *HPB (Oxford)*. 2010;12(1):50-55.
56. Yang T, Zhang J, Lu JH, Yang GS, Wu MC, Yu WF. Risk factors influencing postoperative outcomes of major hepatic resection of hepatocellular carcinoma for patients with underlying liver diseases. *World J Surg*. 2011;35(9):2073-2082.
57. Fisher NC, Mutimer DJ. Central venous cannulation in patients with liver disease and coagulopathy—a prospective audit. *Intensive Care Med*. 1999;25(5):481-485.
58. Bosch J, Abraldes JG, Berzigotti A, García-Pagan JC. The clinical use of HVPG measurements in chronic liver disease. *Nat Rev Gastroenterol Hepatol*. 2009;6(10):573-582.
59. Woolfson J, John P, Kamath B, Ng VL, Ling SC. Measurement of hepatic venous pressure gradient is feasible and safe in children. *J Pediatr Gastroenterol Nutr*. 2013;57(5):634-637.
60. Sleeman D, Namias N, Levi D, et al. Laparoscopic cholecystectomy in cirrhotic patients. *J Am Coll Surg*. 1998;187(4):400-403.
61. da Silveira EB. Outcome of cirrhotic patients undergoing cholecystectomy: applying Bayesian analysis in gastroenterology. *J Gastroenterol Hepatol*. 2006;21(6):958-962.
62. Delis S, Bakoyiannis A, Madariaga J, Bramis J, Tassopoulos N, Derveniz C. Laparoscopic cholecystectomy in cirrhotic patients: the value of MELD score and Child-Pugh classification in predicting outcome. *Surg Endosc*. 2010;24(2):407-412.
63. Carbonell AM, Wolfe LG, DeMaria EJ. Poor outcomes in cirrhosis-associated hernia repair: a nationwide cohort study of 32,033 patients. *Hernia*. 2005;9(4):353-357.
64. Ammar SA. Management of complicated umbilical hernias in cirrhotic patients using permanent mesh: randomized clinical trial. *Hernia*. 2010;14(1):35-38.
65. Castellote J, Xiol X, Cortés-Beut R, Tremosa G, Rodríguez E, Vázquez S. Complications of thoracentesis in cirrhotic patients with pleural effusion. *Rev Esp Enferm Dig*. 2001;93(9):566-575.
66. Xiol X, Castellote J, Cortes-Beut R, Delgado M, Guardiola J, Sesé E. Usefulness and complications of thoracentesis in cirrhotic patients. *Am J Med*. 2001;111(1):67-69.
67. Nielsen SS, Thulstrup AM, Lund L, Vilstrup H, Sørensen HT. Postoperative mortality in patients with liver cirrhosis undergoing transurethral resection of the prostate: a Danish nationwide cohort study. *BJU Int*. 2001;87(3):183-186.
68. Lund L, Jepsen P, Vilstrup H, Sørensen HT. Thirty-day case fatality after nephrectomy in patients with liver cirrhosis—a Danish population-based cohort study. *Scand J Urol Nephrol*. 2003;37(5):433-436. <https://doi.org/10.1080/00365590310006219>
69. Tobin JM, Tanaka KA, Smith CE. Factor concentrates in trauma. *Curr Opin Anaesthesiol*. 2015;28(2):217-226.
70. Maithe SK, Kneuert PJ, Kooby DA, et al. Importance of low preoperative platelet count in selecting patients for resection of

- hepatocellular carcinoma: a multi-institutional analysis. *J Am Coll Surg*. 2011;212(4):638-650.
71. Cinnella G, Pavesi M, De Gasperi A, et al. Standards clinici per il Patient Blood Management e per il management della coagulazione e dell'emostasi nel perioperatorio Position paper della Società Italiana di Anestesia, Analgesia, Rianimazione e Terapia Intensiva (SIAARTI). 2018. <http://www.siaarti.it/SiteAssets/Ricerca/Standards-clinici-per-il-Patient-Blood-Management/SIAARTI%20standard%20PBM.pdf>. Accessed October 5, 2020
 72. Intagliata NM, Argo CK, Stine JG, Lisman T, Caldwell SH, Violi F. Concepts and Controversies in haemostasis and thrombosis associated with liver disease: proceedings of the 7th international coagulation in liver disease conference. *Thromb Haemost*. 2018;118(08):1491-1506. <https://doi.org/10.1055/s-0038-1666861>
 73. Liunbruno G, Bennardello F, Lattanzio A, Piccoli P, Rossetti G. *Raccomandazioni SIMTI sul corretto utilizzo degli emocomponenti e dei plasmi derivati*. Edizioni SIMTI; 2018.
 74. British Committee for Standards in Haematology, Blood Transfusion Task Force. Guidelines for the use of platelet transfusions. *Br J Haematol*. 2003;122(1):10-23.
 75. Violi F, Basili S, Raparelli V, Chowdary P, Gatt A, Burroughs AK. Patients with liver cirrhosis suffer from primary haemostatic defects? Fact or fiction? *J Hepatol*. 2011;55(6):1415-1427.
 76. Basili S, Raparelli V, Napoleone L, et al. Platelet count does not predict bleeding in cirrhotic patients: results from the PRO-LIVER study. *Am J Gastroenterol*. 2018;113(3):368-375.
 77. Hung A, Garcia-Tsao G. Acute kidney injury, but not sepsis, is associated with higher procedure-related bleeding in patients with decompensated cirrhosis. *Liver Int*. 2018;38(8):1437-1441.
 78. Tripodi A, Primignani M, Chantarangkul V, et al. Global hemostasis tests in patients with cirrhosis before and after prophylactic platelet transfusion. *Liver Int*. 2013;33(3):362-367.
 79. Giannini EG, Stravitz RT, Caldwell SH. Correction of hemostatic abnormalities and portal pressure variations in patients with cirrhosis. *Hepatology*. 2014;60(4):1442.
 80. Hidaka H, Kurosaki M, Tanaka H, et al. Lusutrombopag reduces need for platelet transfusion in patients with thrombocytopenia undergoing invasive procedures. *Clin Gastroenterol Hepatol*. 2019;17(6):1192-1200.
 81. Peck-Radosavljevic M, Simon K, Iacobellis A, et al. Lusutrombopag for the treatment of thrombocytopenia in patients with chronic liver disease undergoing invasive procedures (L-PLUS 2). *Hepatology*. 2019;70(4):1336-1348.
 82. Terrault N, Chen YC, Izumi N, et al. Avatrombopag before procedures reduces need for platelet transfusion in patients with chronic liver disease and thrombocytopenia. *Gastroenterology*. 2018;155(3):705-718.
 83. Loffredo L, Violi F. Thrombopoietin receptor agonists and risk of portal vein thrombosis in patients with liver disease and thrombocytopenia: a meta-analysis. *Dig Liver Dis*. 2019;51(1):24-27.
 84. Afdhal NH, Giannini EG, Tayyab G, et al. Eltrombopag before procedures in patients with cirrhosis and thrombocytopenia. *N Engl J Med*. 2012;367(8):716-724.
 85. Committee for Medicinal Products for Human Use (CHMP). Assessment report – Lusutrombopag Shionogi. EMA/CHMP/817852/2018. https://www.ema.europa.eu/en/documents/assessment-report/lusutrombopag-shionogi-epar-public-assessment-report_en.pdf. Accessed October 5, 2020
 86. Fortea JJ, Puente Á, Ezcurra I, et al. Management of haemostatic alterations and associated disorders in cirrhosis in Spain: a national survey. *Dig Liver Dis*. 2019;51(1):95-103.
 87. Desborough MJ, Hockley B, Sekhar M, et al. Patterns of blood component use in cirrhosis: a nationwide study. *Liver Int*. 2016;36(4):522-529.

How to cite this article: Alvaro D, Caporaso N, Giannini EG, et al; The PREBRiC (Procedure-Related Bleeding Risk in Cirrhosis) group. Procedure-related bleeding risk in patients with cirrhosis and severe thrombocytopenia. *Eur J Clin Invest*. 2021;51:e13508. <https://doi.org/10.1111/eci.13508>