

Review

Salvia Species: Biotechnological Strategies Applied to In Vitro Cultures for the Controlled Production of Bioactive Diterpenoids

Roméo Arago Dougué Kentsop ^{1,†} , Poonam Devi ², Andrea Copetta ³ , Barbara Ruffoni ³ , Valentina Parisi ⁴, Angela Bisio ^{2,*}  and Valeria Iobbi ^{2,*} 

¹ Institute of Agricultural Biology and Biotechnology, National Research Council, Via Bassini 15, 20133 Milan, Italy; romeo.dougue@ibba.cnr.it

² Department of Pharmacy, University of Genova, Viale Cembrano 4, 16148 Genova, Italy; poonam.devi@edu.unige.it

³ Consiglio per la Ricerca e la Sperimentazione in Agricoltura—CREA Centro di Ricerca Orticoltura e Florovivaismo, 18038 Sanremo, Italy; andrea.copetta@crea.gov.it (A.C.); barbara.ruffoni@crea.gov.it (B.R.)

⁴ Department of Pharmacy, University of Salerno, via Giovanni Paolo II 132, 84084 Fisciano, Italy; vparisi@unisa.it

* Correspondence: angela.bisio@unige.it (A.B.); valeria.iobbi@edu.unige.it (V.I.)

† This paper is a part of the Ph.D. Thesis of Roméo Arago Dougué Kentsop, presented at the University of Genoa, Genoa, Italy.

Abstract: Plant secondary metabolites have great applications in the nutritional and cosmetic aspects of human health. Terpenes, and in particular bioactive diterpenoids, represent an important group of compounds found in *Salvia* species. Their production in plants is often limited, and chemical synthesis is often not economically feasible. Biotechnological approaches using plant cell and tissue cultures can be routinely established under sterile conditions from explants for biomass production and the extraction of secondary metabolites. The biosynthesis and accumulation of bioactive diterpenoids in vitro in *Salvia* cells and tissues can be enhanced by strain improvement, techniques to select high-producing cell lines, optimisation of the growth medium, use of specific precursors or elicitors, induction of their release into the culture medium, and the overexpression of genes. This review analyses the biotechnological techniques applied to the in vitro culture of *Salvia* cells and tissues to enhance the production and accumulation of bioactive diterpenoids and summarises their biological activities.

Keywords: diterpenoids; *Salvia*; elicitation; in vitro culture; hairy root; cell culture; biological activity



Citation: Dougué Kentsop, R.A.; Devi, P.; Copetta, A.; Ruffoni, B.; Parisi, V.; Bisio, A.; Iobbi, V. *Salvia* Species: Biotechnological Strategies Applied to In Vitro Cultures for the Controlled Production of Bioactive Diterpenoids. *Agronomy* **2024**, *14*, 835. <https://doi.org/10.3390/agronomy14040835>

Academic Editor: Kirsten Brandt

Received: 20 March 2024

Revised: 9 April 2024

Accepted: 16 April 2024

Published: 17 April 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Salvia L. is a large genus of the Lamiaceae, with about 980 species. It is mainly distributed in Mesoamerica/South America, the Mediterranean region, and Southwest and East Asia [1–5]. *Salvia* species have long been known for their wide variety of medicinal uses in folk medicine for pain relief, protection against oxidative stress, free radical damage, angiogenesis, inflammation, and bacterial and virus infection [6–10]. *Salvia* are cultivated for their aromatic properties and abundance of secondary metabolites, which are used to produce food additives and essential oils, pharmaceuticals, dyes, cosmetics, and bio-cides [8,11]. Due to their widespread ethnic use, some *Salvia* species (*S. sclarea*, *S. officinalis*, and *S. fruticosa*) are recognised as safe in the USA [12], and their drugs are included in the European Pharmacopoeia VI. From a phytochemical point of view, *Salvia* spp. have a complex chemical composition with terpenoids [7,13–16] and polyphenols [17–20] being the main compounds. By the beginning of 2011, a total of about 773 compounds had been isolated from 134 *Salvia* species and classified into seven main groups: sesquiterpenoids, diterpenoids, sesterterpenoids, triterpenoids, steroids, polyphenols, and others. According to their structure, diterpenoids can further be divided into a number of groups [8,21]. The most

representative are abietane [22–24], clerodane [25–28], pimarane [29], icetexane [30–32], and labdane [13]. Extensive research into biotechnological approaches to the production of *Salvia* phytochemicals has been driven by the increasing demand for sage products and the environmental, ecological, and climatic constraints of producing sage metabolites from field-grown plants [33]. Growing demand has led to the overexploitation of natural habitats, and complex climate-related stressors can affect many wild *Salvia* species [34]. Alternative and renewable sources of *Salvia* biomass are urgently needed to prevent ecological crisis. In vitro plant culture technology is a potential approach for the continuous production of secondary metabolites in *Salvia* species under controlled conditions. In vitro systems are environmentally friendly and can be grown independently of natural plant populations, environmental conditions, latitude, climatic variations, or seasonal changes. In vitro plant tissues and cell culture methods may represent an alternative strategy for the production of highly valuable plant metabolites, and could even be adopted by the pharmaceutical industry for the development of new drugs and formulations, the production of pharmaceuticals and cosmetics, as food additives in the food industry, and for crop protection in agriculture [35,36]. Biotechnological techniques have greatly facilitated plant propagation and the production of some important bioactive compounds from the genus *Salvia*. Various in vitro systems such as shoots, cell culture, and hairy roots are used to produce bioactive compounds [33,37–44]. However, many issues remain to be addressed before large-scale biotechnological production based on *Salvia* in vitro systems can be developed [41].

This review, which originated from the first author's PhD thesis [45], focuses on the main diterpenoids obtained from in vitro cultures of *Salvia* species. The aim of the present review is to provide an overview of promising biotechnological strategies for the precise control of growth conditions and the optimisation of growth regulators, nutrient supply, and elicitation strategies to stimulate diterpenoid biosynthesis in in vitro *Salvia* species. Furthermore, genetic engineering techniques to modify metabolic pathways to enhance the production of specific bioactive diterpenoids have also been reported. To the best of our knowledge, this is the first review article dealing with the in vitro production of different classes of diterpenoids in different *Salvia* species.

2. Search Strategy

The present paper provides an overview of the most important results reported in the literature on the in vitro production of bioactive diterpenes from cell and tissue cultures of *Salvia* species. A systematic search was conducted in English language databases (Medline (PubMed), Web of Science, and Scopus) from inception to January 2024, with no restriction on publication date. In addition, report papers and reviews were manually retrieved, and traced references in the included literature were tracked down using the references found to retrieve additional material retrospectively. The articles used in this review were selected from journals with a high impact factor, based on their scientific relevance and the reliability of their data. The search was carried out independently by three researchers and then cross-checked. Keywords and phrases used in the search included the following: "*Salvia*", "diterpenoids", "bioactive diterpenes in *Salvia*", "biological activity", "elicitation", "biotic elicitors", "abiotic elicitors", "combination of elicitors", "in vitro cultures", "hairy root", "cell culture", "growth conditions", "in situ adsorption", "medium renewal", and "gene overexpression". In total, 19 *Salvia* species were considered. For a better understanding of the studies presented in this review, tables in the Supplementary Materials (Tables S1–S9) were prepared to show the main results related to the influence of the elicitation on producing and accumulating specific bioactive diterpenes.

3. Structure and Biological Activity of the Main Diterpenoids Obtained from In Vitro Cultures of *Salvia* Species

Diterpenoids [16,46–62] are composed of four isoprene units and 20 carbon atoms in their chemical structures. The various bioactive diterpenoids found in the in vitro culture

of tissues or cells in *Salvia* species are detailed in Table 1, and the structures of the major bioactive diterpenoids found in in vitro *Salvia* species are reported in Figure 1.




Figure 1. Bioactive diterpenoids obtained from in vitro cell and tissue cultures of *Salvia* species. Reproduced from [45].

Table 1. Diterpenoids produced by the in vitro culture of various *Salvia* species. Reproduced from [45].

Compound Number	Name	<i>Salvia</i> Species	In Vitro Culture	References
1	tanshinone I	<i>S. miltiorrhiza</i>	HR	[63]
			CC	[64]
2	tanshinone IIA	<i>S. castanea</i>	HR	[65]
			HR	[63]
			CC	[64]
3	cryptotanshinone	<i>S. miltiorrhiza</i>	HR	[63]
			CC	[64]
			HR	[65]
4	dihydrotanshinone I	<i>S. miltiorrhiza</i>	HR	[63]
			HR	[65]
5	salvipisone	<i>S. sclarea</i>	HR	[66]
6	aethiopinone	<i>S. sclarea</i>	HR	[66]
7	1-oxoaethiopinone	<i>S. sclarea</i>	HR	[66]
8	ferruginol	<i>S. sclarea</i>	HR	[66]
9	carnosic acid	<i>S. miltiorrhiza</i>	HR	[63]
			CC	[67]
			MP	[37]
10	carnosol	<i>S. officinalis</i>	C	[67]
			CC	[67]
			MP	[37]
11	taxodone	<i>S. austriaca</i>	HR	[68]
12	taxodione	<i>S. austriaca</i>	HR	[68]
13	15-deoxy-fuerstione	<i>S. austriaca</i>	HR	[68]
14	demethylfruticuline A	<i>S. corrugata</i>	RS	[38]

Table 1. Cont.

Compound Number	Name	Salvia Species	In Vitro Culture	References
15	fruticuline A	<i>S. corrugata</i>	MP	[38]
			apex	[38]
			leaves	[38]
			RS	[38]
			apex	[38]
			leaves	[38]

HR: hairy roots, CC: cell culture, C: callus, MP: micropropagated plants, RS: regenerated shoots.

3.1. Tanshinones: Tanshinone I, Tanshinone IIA, Cryptotanshinone, Dihydrotanshinone I

Tanshinones are widely distributed in the genus *Salvia* [7,69–71]. The dried root of *S. miltiorrhiza* is commonly used in Traditional Chinese Medicine (TCM) to improve body function (e.g., stimulate circulation and improve blood flow) [72]. Several hydrophobic bioactive diterpenoids have been isolated from this drug [9,70,72–79]. Since the 1990s, several researchers have exploited the in vitro culture of *S. miltiorrhiza* including the hairy roots and cell culture to produce tanshinones. Due to their remarkable activities in the clinical treatment of cardiovascular diseases, these compounds have received much attention. The roots of *S. castanea*, *S. tomentosa*, and *S. przewalskii* have been used as a substitute for *S. miltiorrhiza* roots because of their tanshinone content [80–82]. Several years later, roots of micropropagated and in vitro regenerated *S. przewalskii* plants were assayed for tanshinone I (T-I) and tanshinone IIA (T-IIA). The results showed that the highest tanshinone levels, 3.8 mg/g dry weight (DW) of TI and 7.6 mg/g DW of TIIA, were produced by 2-year-old roots in in vitro regenerated plants grown in the field and harvested at the flowering stage [83]. Tanshinones have also been found in the roots of *S. abrotanoides* (ex *Perovskia abrotanoides*) [84], and the amounts are increased by biotic and abiotic elicitors [85]. Nowadays, these compounds have been found to possess a wide range of pharmacological activities such as antibacterial, antioxidant, anti-inflammatory, and antitumour properties [86].

3.2. Abietane Diterpenoids: Aethiopinone, Salvipisone, Ferruginol, and 1-Oxo-aethiopinone

The rearranged abietane diterpenoids salvipisone (5) and aethiopinone (6) as well as ferruginol have been extracted from the roots of *S. aethiopsis* [87–89]. These compounds are mainly present in the roots of some *Salvia* species and have also been isolated from several species such as *S. argentea* [90], *S. candidissima* [91], *S. ceratophylla* [92], and from the aerial parts of *S. cyanescens* [93]. All of these diterpenoids as well as the rearranged abietane 1-oxo-aethiopinone (7) have been isolated from the roots of *S. sclarea* [94]. Only *S. sclarea* has been investigated for producing these abietane diterpenoids using hairy root transformation [95,96]. As previously described, ferruginol (8), salvipisone, aethiopinone, and 1-oxo-aethiopinone were produced by the in vitro hairy root transformation of *S. sclarea* using *Agrobacterium rhizogenes* strain LBA 9402 [95]. Ferruginol has recently been purified from the hairy roots of *S. corrugata* [97]. These compounds have been reported to have various biological activities. Among them, ferruginol exhibited potent anti-acanthamoeba [98] and in vitro antiplasmodial activities [99]. Aethiopinone has been shown to have analgesic and anti-inflammatory properties [100]. It has also been shown to be cytotoxic to several solid tumour cell lines. The most potent effect was on the human melanoma A375 cell line (IC₅₀ 11.4 µM) [66]. Salvipisone and aethiopinone have shown antibacterial activity against selected methicillin-susceptible and methicillin-resistant *S. aureus* and *S. epidermidis* strains [101–103]. These compounds exhibited relatively high cytotoxicity against HL-60 and NALM-6 leukaemia cells (IC₅₀ range 0.6–7.7 µg/mL, corresponding to 2.0–24.7 µM), whereas 1-oxo-aethiopinone and ferruginol were less active [104].

3.3. Abietane Diterpenoids: Carnosic Acid and Carnosol

Carnosic acid (**9**) is one of the most studied phenolic diterpenoids [105,106]. It is important in the food, cosmetic, and pharmaceutical industries because of its potent antioxidant, anti-inflammatory, and anticancer properties [107–115]. Carnosic acid and carnosol (**10**) have been isolated from various *Salvia* species [87,116]. Carnosic acid is mainly found in the aerial parts of the plant, and its biosynthesis in the leaves has been extensively studied in *S. fruticosa* and *S. rosmarinus* [106,117,118]. The biological activity of carnosic acid and carnosol has been the subject of extensive investigation [119]. Carnosic acid and carnosol exhibited anti-inflammatory activities by acting on mPGES-1 and 5-LO, resulting in suppressing pro-inflammatory eicosanoid formation [114]. Later on, these diterpenoids from the leaves of *S. officinalis* were found to have antioxidant and antibacterial activities [108]. Recently, carnosic acid was shown to suppress the development of oral squamous cell carcinoma via the mitochondrial apoptotic pathway [120] and to prevent the biofilm formation of methicillin-resistant *Staphylococcus aureus* (MRSA) [121]. Recent studies have demonstrated its potential as a preventive agent against soft rot caused by *Pectobacterium carotovorum* subsp. *carotovorum* on slices of potato tubers [122].

3.4. Abietane Diterpenoids: Taxodone, Taxodione, and 15-Deoxyfuerstione

Taxodone (**11**), taxodione (**12**), and 15-deoxyfuerstione (**13**) were isolated from the roots of *S. spinosa* [123] and the hairy roots of *S. austriaca* [68]. Taxodione showed activity against human acetylcholinesterase and cytotoxic activity against the A549 cell line [124]. This compound was found to be active against *Trypanosoma brucei rhodesiense* (IC₅₀ = 0.05 µM with high selectivity, SI = 38) and inhibited the growth of *Plasmodium falciparum* and *T. cruzi* by 50% at concentrations of 1.9 and 7.1 µM, respectively (SI values of 1.0 and 0.27) [68]. Taxodione from the roots of *S. deserta* was found to be leishmanicidal with an IC₅₀ value of 1.46 µM (0.46 mg/L) against *Leishmania donovani* [125]. Taxodione from *S. chorassanica* exerted a protective effect in ischaemic injury induced by serum/glucose deprivation and the putative role of apoptosis as an underlying mechanism [126]. Taxodone and 15-deoxy-fuerstione are potent antimicrobial agents. In addition to direct biocidal/biostatic activity, they also interfered with the primary virulence factors/mechanisms of *S. aureus* and *Candida albicans*. In addition, both diterpenoids significantly inhibited microbial adhesion and biofilm formation when used at sub-inhibitory concentrations [127]. 15-Deoxyfuerstione from the roots of *S. lachnocalyx* showed a cytotoxic activity against MOLT-4, HT-29 and MCF-7 cells and significant anticancer activity with IC₅₀ values in the range of 0.54–11.82 mg/mL [123].

3.5. Icetexane Diterpenoids: Demethylfruticuline A and Fruticuline A

Demethylfruticuline A (**14**) and fruticuline A (**15**) have been reported as the main constituents of the aerial part of *S. corrugata* [30]. Fruticuline A is an icetexane diterpenoid first isolated from *S. fruticulosa* [128], and later from *S. arizonica* [19], *S. corrugata* [30], and *S. lachnostachys* [129]. As previously described for the establishment of an in vitro culture of *S. corrugata*, analysis of the regenerated shoots showed the presence of both icetexanes. The yield of fruticuline A was higher in the methanolic extract than in those of fresh leaves and fresh shoot tips. However, only fruticuline A was found in micropropagated plants, whereas trace amounts of both diterpenoids were found in callus [38]. The biosynthetic pathway of demethylfruticuline A, fruticuline A has not been reported. However, biosynthetic relationships have shown that the icetexane skeleton originates from a rearrangement of the more common abietane skeleton, resulting in a 6-7-6 tricyclic scaffold with the systematic name 9(10/20)-abeo-abietane [23]. Consistent with this hypothesis, the majority of icetexane natural products discovered to date have been found in plant species that also produce abietane diterpenoids as secondary metabolites [31]. The icetexanes discovered to date vary widely in the degree of oxygenation and oxidation in each ring, resulting in a range of structures and biological activities [32]. Fruticuline A and demethylfruticuline A have shown potent antibacterial activity against Gram-positive bacteria [130,131], and

antioxidant activity [132]. Fruticuline A has also been reported to have anti-inflammatory and analgesic properties [133]. Demethylfruticuline A induces anoikis, a type of apoptosis induced in mammalian cells by the loss of cell adhesion mediated by CD36 [134], and also causes apoptosis by the induction of reactive oxygen species in mitochondria [135]. Recently, fruticuline A and demethylfruticuline A have shown remarkable binding affinity to CDK-2 compared to known CDK-2 inhibitors, leading to the consideration of these compounds as excellent natural CDK-2 inhibitors [136].

4. Strategies to Increase the In Vitro Production of Diterpenoids

Several parameters should be considered to control the optimal in vitro production of bioactive secondary metabolites from plants. Medium composition (sucrose level, exogenous growth hormone, type of nitrogen source, and their relative amounts) and growth conditions (light, temperature, and the presence of chemicals) can influence the growth, total biomass yield, and secondary metabolite production [137,138]. Specific treatments of the in vitro culture including precursor feeding and elicitor application can be used to increase plant metabolite production and improve its qualitative value as an ingredient in the pharmaceutical industry [139,140]. All of these approaches have opened up a new area of research that could have important economic benefits. For *Salvia* species, the differentiation or not of tissue culture, the chemical composition of the culture medium, the use of elicitors, the interaction with microorganisms (bacteria or fungi), the reduction in extracellular secondary metabolites by absorption, and transgenic culture with genes involved in the biosynthetic pathway are some in vitro strategies to increase the production of bioactive diterpenoids.

4.1. Medium Composition and Effect of Nutrients (Mineral, Hormone, and Sucrose Composition)

The in vitro growth of plant organs, tissues, and cells takes place in an appropriate medium that contains all of the necessary elements (salt composition, percentage of sucrose, and the presence of plant growth regulators). These elements have an influence on the accumulation of biomass and the formation of secondary metabolites. External factors such as carbon source, nitrogen source, growth regulators, medium pH, temperature, light, and oxygen are considered to easily regulate the expression of plant secondary metabolite pathways [141,142].

4.1.1. Effect of Salts

The most commonly used media are Murashige and Skoog (MS) [143], with very high concentrations of nitrate (NO_3^-), potassium (K^+), and ammonia (NH_4^+), Gamborg (B5) [144] and 67-V [145], with lower levels of inorganic nutrients than MS. Salt composition can affect biomass growth and secondary metabolite production. In general, high concentrations of ammonium ions inhibit the formation of secondary metabolites, while lowering the ammonium nitrogen increases it [146]. Zhi et al. [63] showed that the accumulation of diterpenoids in the hairy roots of *S. miltiorrhiza* was higher in the absence of ammonium nitrate (about 220 mg dry roots and 19 mg diterpenoids/g DW) than in its presence (about 140 mg dry roots and 7 mg diterpenoids/g DW). In the *Ti*-transformed cell suspension cultures, the B5 medium supported the best growth while the 67-V medium promoted tanshinone production [64]. Compared to the control, changing the KH_2PO_4 concentration could promote adventitious growth, but a high KH_2PO_4 concentration inhibited T-IIA biosynthesis in adventitious roots of *S. miltiorrhiza* [147].

4.1.2. Effect of Hormones

Cytokinin types and concentrations have different effects on different metabolites. Absciscic acid (ABA) and thidiazuron (TDZ) promote the increase in T-I and cryptotanshinone (CT) 5-fold and 7.5-fold, respectively, over the control in *S. miltiorrhiza* hairy root cultures [148]. In *S. officinalis*, the content of carnosol reached the highest value of 30.5 ± 0.42 mg/g DW in suspension cultures supplemented with 1.5 mg/L zeatine (ZEA),

and carnosic acid reached 12.7 ± 2.96 mg/g DW in callus cultures supplemented with 1.5 mg/L 6-benzylaminopurine (BA). However, the amount of these compounds was at a trace level in the suspension cultures supplemented with 0.5 mg/L of kinetin (KIN) [67].

4.1.3. Effect of Sucrose

The initial concentration of sucrose, which is the main carbon source in plant culture, has a significant effect on the growth of transformed cell cultures of *S. miltiorrhiza*. The fastest growth was observed in medium containing 30 g/L sucrose; it was significantly reduced at 60 g/L sucrose, and completely inhibited at 100 g/L sucrose [64]. The level of sucrose affects the productivity of secondary metabolite accumulating cultures [149]. In Ti-transformed *S. miltiorrhiza* cell suspension cultures, tanshinone production was better in the 30 g/L sucrose medium than in the 20 g/L or 40 g/L sucrose medium. However, a much higher tanshinone production was observed in the 100 g/L sucrose medium, although the cell growth was almost completely inhibited under these conditions [64].

4.1.4. Effect of Environment Factors

Light plays an important role in promoting plant growth and inducing or regulating plant metabolism [150–153]. On the other hand, light has an inhibitory effect on the accumulation of secondary metabolites such as nicotine and shikonin [154]. The green cell aggregates of *S. miltiorrhiza* transferred to the fresh B5 medium containing 4 g/L yeast extract and cultivated in darkness produced more tanshinones than their counterparts cultivated under continuous illumination, suggesting the inhibitory effect of light on tanshinone production in the cell cultures [64].

4.2. Elicitation

Plants and in vitro cultured plant cells show physiological and morphological responses to microbial, physical, or chemical factors, which are known as “elicitors” [155]. The term elicitor was originally used to describe molecules capable of inducing the production of phytoalexins. It is now defined as a substance that, when applied in small amounts to a living system, can induce or enhance the biosynthesis of specific compounds that are critical for plant adaptation to stress conditions [156]. Table 2 classifies elicitors according to their nature and may include abiotic elicitors such as metal ions and inorganic compounds as well as biotic elicitors from fungi, bacteria, viruses, or herbivores, plant cell wall components, and chemicals released by plants at the site of infection when attacked by pathogens or herbivores [157]. Elicitation is a widely used method that aims to enhance the production of secondary metabolites and has been described for many plant species. The discovery of elicitors has opened up a new approach to secondary metabolite production. All culture types (cell, callus, root, hairy root, whole plant, shoots, and seedling) can be exposed to elicitor treatments. Methyl jasmonate (MJ), jasmonic acid (JA), and salicylic acid (SA) are the most commonly used elicitors. In addition, polysaccharides such as pectin, dextran, chitin, chitosan, and alginate are often used as elicitors to induce secondary metabolism in plant cell and tissue cultures [158]. For an appropriate effect, several parameters such as elicitor concentration and selectivity, duration of elicitor exposure, age of culture, cell line, growth regulation, nutrient composition, and quality of the cell wall materials are also important factors influencing the successful production of secondary metabolites [159].

Table 2. Elicitors classification based on their nature. Adapted from Naik and Al-Khayri [160].

Biotic Elicitors	Abiotic Elicitors			
	Physical	Chemical	Hormones	Plant Signal Compounds
Polysaccharide	UV radiation	Heavy metals	ABA	Jasmonic acid
Yeast extract	Osmotic stress	Mineral salts	TDZ	Salicylic acid
Fungi	Salinity	Gaseous toxins		
Bacteria	Drought			
	Thermal stress			

The activation of the biosynthesis of the plant product requires the perception of an extracellular or intracellular signal by a receptor on the surface of the plasma membrane. Elicitor perception initiates a signal transduction cascade, leading to the activation or de novo biosynthesis of transcription factors. These in turn directly regulate the expression of biosynthetic genes involved in secondary metabolism [161].

4.2.1. Biotic Elicitors

Microorganism—Roots Interaction

The elicitor-induced production of secondary metabolites in culture has been tested in a variety of microorganisms (viruses, bacteria, algae, and fungi), not necessarily pathogens. Rhizosphere microbes are best known to act as biotic elicitors that can induce the synthesis of secondary metabolites in plants [157]. These microorganisms are mainly endophytes (microorganisms that colonise the inside of host plants without causing any apparent disease) and have been used by some researchers in co-culture with their host plant tissues as efficient elicitors leading to the production of secondary metabolites. In some cases, these microorganisms have a direct interaction with the plant tissues. The amount of inoculum is important for inducing the elicitation effect. However, not all endophytes can co-culture with their host plant tissues for a long time, depending on the toxicity of the fungal isolate. In most cases, these fungi have been produced as elicitors with their toxicity removed, which can also stimulate the secondary metabolism of the host plants [162]. The endophyte has a symbiotic relationship with the plants, which may have a long evolutionary and interactive communication. Table S1 (Supplementary Materials) reports the effect of microorganism interaction and their extracts on diterpenoid production in *in vitro* *Salvia* hairy root cultures.

Actinomycetes are an abundant and widespread group of soil microbes, constituting approximately 10 to 50% of the soil microbial community. Tyc et al. [163] and Adegbeye et al. [164] and others have reported them to be important producers of secondary metabolites [165]. These microorganisms have been widely studied for their ability to modulate the relationships between plants and biotic/abiotic stresses, often producing valuable secondary metabolites that can affect host physiology [166]. A soil actinomycete strain, designated *Streptomyces pactum* Act12, has been isolated from the drought, low temperature, and high latitude environment of the Qinghai-Tibetan Plateau [167]. Most *Streptomyces* are efficient rhizosphere and rhizoplane colonisers. They can also be endophytes, colonising the internal tissues of host plants [168]. Yan et al. [169] showed that hairy roots of *S. miltiorrhiza* treated with 2% and 4% *Streptomyces pactum* Act12 for 14 days strongly promoted the synthesis of tanshinones, most significantly increasing the concentration of CT 13.21 and 33.63 times more than that of the control at 2% and 4% *Streptomyces* treatment as well as 8.42 and 15.31 for DT-I, 11.77 and 10.36 for T-IIA, and 5.58 and 5.67 for T-I concentrations, respectively. The stimulatory effect of ACT12 on T-I was less than on some of the other tanshinones because the concentrations of the four substances were significantly increased. The total tanshinone concentrations in the hairy roots of *S. miltiorrhiza* were 9.19 and 12.61 higher in the 2% and 4% ACT12 treatments, respectively.

Plant growth promoting rhizobacteria (PGPR) are rhizosphere bacteria that can colonise the root system (rhizosphere, rhizoplane, or roots) of the plants. They are beneficial to plant growth by facilitating nutrient uptake, producing and releasing growth-promoting substances such as phytohormones, and providing plant protection [170,171]. *Bacillus cereus* is one of the most common rhizobacterial species that has been shown to increase plant resistance to bacterial and fungal pathogens [170,172]. Research by Wu and co-workers [173,174] showed that live *B. cereus* cells inoculated into the *S. miltiorrhiza* hairy root culture on day 0 dramatically stimulated the total root tanshinone accumulation but suppressed hairy root growth. The culture inoculated with 0.2% (OD value of 0.5) bacteria on day 0 reached the highest tanshinone content of 2.67 mg/g DW and the highest volumetric yield of 10.4 mg/L (on day 28), which were 13.5 and 7.6 times higher than those of the control at 0.20 mg/g DW and 1.40 mg/L, respectively [174]. A few years later, the tanshinone accumulation was

most dramatically enhanced by a bacterial concentration of 2.5% (with an OD value of 0.5) inoculated on day 0 for 7 days to 2.78 mg/g DW, which was about 18-fold higher than in the control culture at 0.15 mg/g DW. In addition, the highest volumetric tanshinone yield of 22.4 mg/L, which was approximately 12-fold higher than the control yield of 1.82 mg/L, was achieved with the same bacterial concentration inoculated on day 18 [173].

Yan et al. [175] described that *Pseudomonas brassicacearum* sub sp. *neaurantiaca* (B1), isolated from both the phloem and xylem of healthy *S. miltiorrhiza*, inoculated into *S. miltiorrhiza* hairy root culture at 0.025% (OD value of 0.5) on day 21 and maintained in co-culture for 9 days, promoted the biomass growth. In addition, it promoted the increase in the total tanshinone content in hairy roots, reaching a 3.7-fold increase compared to the control, and most obviously increased the content of dihydrotanshinone I (DT-I) and CT content (19.2-fold and 11.3-fold, respectively) compared to the control. Consequently, hairy roots of *S. miltiorrhiza* have inhibitory effects on the growth of endophytic bacteria B1, which is mainly due to the antibacterial effect of tanshinone substances in the hairy roots.

Zhai et al. [176] investigated the effects of the live endophytic fungus *Chaetomium globosum* D38 on tanshinone biosynthesis in *S. miltiorrhiza* hairy roots and showed that administration of D38 on day 18 significantly increased the levels of DT-I and CT by 8-fold and 14.9-fold, respectively, compared to the control.

Microorganism Extracts and Constituents

Biotic elicitors produced by pathogens have been mainly used in *Salvia* species to induce the plant defence responses and secondary metabolite production.

Trichoderma species control fungal pathogens both by microbial antagonism and by inducing local and systemic responses [177]. The endophytic fungus *Trichoderma atroviride* D16 from the root of *S. miltiorrhiza* was first reported to produce T-I and T-IIA in mycological medium [178]. A few years later, Ming et al. [179] showed that both the mycelium extract (EM) and polysaccharide fraction (PSF) from *Trichoderma atroviride* D16 treatment were responsible for promoting hairy root growth and causing a significant change in the abundance of the four tanshinones. Among the four tanshinone species, DT-I and CT were most dramatically stimulated by EM and PSF on day 18. The content of DT-I in hairy roots treated with 300 mg/L EM was ~35-fold higher than the control (1.338 mg/g DW versus 0.039 mg/g DW), and the content of CT treated with 150 mg/L EM was almost 83-fold higher than that of the control (3.061 mg/g DW versus 0.037 mg/g DW). Similarly, the levels of DT-I and CT in hairy roots treated with 180 mg/L PSF were ~23-fold (1.216 mg/g DW versus 0.052 mg/g DW) and ~66-fold (3.496 mg/g DW versus 0.053 mg/g DW) higher, respectively, than the control. Furthermore, under the influence of *C. globosum* D38 EM, the content of DT-I and CT reached the highest levels. These were 21-fold and 19.8-fold higher than the control group at doses of 60 mg/L and 90 mg/L, respectively [176].

Chitosan and chitin are structural components of cell walls found in many fungi. However, chitosan is a hydrophilic biopolymer obtained by the *N*-deacetylation of chitin and can be used as an antimicrobial agent [180]. Zhao et al. [181] showed that the application of 100 mg/L chitosan in *S. miltiorrhiza* cell culture induced the production of T-I 3.4-fold (0.27 mg/g versus 80 µg/g of the control). The authors also found that its stimulating effect on tanshinone accumulation (about 6-fold) was stronger than SA, MJ, and sorbitol, but much weaker than Ag⁺, Cd, and YE.

Bacterial extracts are biological mixtures prepared from autoclaved and centrifuged microorganism cultures without identification of the active compounds [182]. The *Bacillus cereus* bacterial extract slightly enhanced both the hairy root growth and tanshinone biosynthesis when it was fed to the *S. miltiorrhiza* hairy root culture growing on MS medium on day 21. Wu et al. [174] described that the bacterial extract of 100 mg/mL applied on day 0 and harvested 28 days after the application increased the root weight by almost 50% and the TT content of roots increased by about 2-fold compared to the control. Furthermore, Zhao et al. [173] showed that the 10-fold higher concentration of bacterial extract (1 g/L) applied on day 0 and harvested 7 days later dramatically reduced the hairy root biomass by

5.21 g DW/L and 7.54 g DW/L compared to the control at 12.2 g DW/L and significantly increased the tanshinone content by about 12-fold (2.04 mg/g DW versus 0.17 mg/g DW in the control).

In addition, coronatine (Cor), an analogue of methyl jasmonic acid, a polyketide effector molecule produced by *Pseudomonas syringae* pv *tomato* strain DC3000 (Pst DC3000), bound the same receptor and acted as a structural agonist of JA-Ile [183]. It was first described as an elicitor by Weiler et al. [184]. Vaccaro et al. [185] showed that 0.1 μ M Cor, applied to *S. sclarea* hairy root cultures for 28 days, allowed the extraction of 103.32 ± 2.10 mg/L aethiopinone, corresponding to approximately a 24-fold increase over the basal content of the control hairy roots (4.40 ± 0.13 mg/L). This level was higher than the 16-fold increase (73.29 ± 0.11 mg/L) induced by elicitation with MJ for the same elicitation time, and higher than the final yield obtained after 7 days with this elicitor (24.08 ± 0.79 mg/L). Interestingly, Cor increased the content of carnosic acid, with the maximum yield obtained after 28 days of elicitation (36.75 ± 2.20 mg/L), representing an approximately 18-fold increase over the basal content of the control hairy roots (Table S2, Supplementary Materials).

For decades, scientists have been using yeast extract as one of the biotic elicitors. Despite limited knowledge of the composition and mechanism of action of yeast, autoclaved solutions providing cell wall fragments have been widely used as elicitors to enhance plant secondary metabolite production, mainly in plant cell or hairy root cultures [182]. Treatments of transformed and nontransformed *S. miltiorrhiza* cell cultures with a yeast extract mainly resulted in an inhibition of biomass accumulation (Table S3, Supplementary Materials). Nevertheless, Ti transformed cells cultured in fresh 6,7-V medium containing 20 g/L sucrose with 4 g/L yeast extract for 8 days stimulated the production of the total tanshinone (CT and T-IIA) productivity of 22.2 mg/L compared to trace amounts in the control cultured in fresh B5 medium [64]. When cultured in a MS-NH₄ medium (MS without ammonium nitrate, containing 30 g/L sucrose) yeast elicitor (4 g/L) for 18 days, this production increased from trace levels in the control to 12.23 mg/L [186]. Treatment of an 18-day-old nontransformed *S. miltiorrhiza* cell culture with yeast extract at 100 mg/L for 7 days reduced biomass production by no more than 50% (5.1–5.5 g/L versus 8.9 g/L), but increased the total tanshinone content to 2.30 mg/g, about 11.5 times that of the control (0.20 mg/g), and more drastically, the CT content was about 2.01 μ g/g, which was 34 times that of the control (60 μ g/g) [181]. This last result is in agreement with that of the normal cell suspensions of *S. miltiorrhiza* in which CT production was stimulated only after cell growth had been suppressed [187]. However, when 4-day-old Ti transformed cells were treated with 0.1% (v/v) YE, the biomass accumulation decreased compared to the control (13.1 versus 14.4 g DW/L at day 5) while the CT production increased significantly (11.5 versus 0 mg/L) [188].

Table S4 (Supplementary Materials) illustrates the effects of the yeast extract on diterpenoid production in *in vitro* *Salvia* hairy root cultures. In contrast to the cell cultures, applying the yeast extract to the hairy roots did not only inhibit the biomass production. Treatment of 18-day-old hairy roots of *S. miltiorrhiza* for 9 days with YE 100 mg/L increased the volumetric total tanshinone (CT, T-I, and T-IIA) of 7.62 mg/L about 4.3-fold over the control [189], and a treatment for 4 days increased the total tanshinone by about ~2.2 mg/g DW, which was 3.1-fold over the control [190]. However, the treatment with YE 25 mg/L increased this amount to 9.92 mg/L, approximately 5-fold over the control [191]. The total tanshinone (CT, T-I, T-IIA) increased significantly when 18-day-old hairy roots were treated with 100 mg/L for 12 days, reaching 13.7 mg/L, about 3.8 times the control [192]. Yang et al. [193] investigated the production of tanshinones when *S. miltiorrhiza* hairy roots were treated with 200 mg/L YE and showed that CT was the most produced diterpenoid. Chen et al. [194] reported that the intracellular content of CT increased from 0.001% to as much as 0.096% of DW.

Treatment of *S. castanea f. tomentosa* with an optimum concentration of 200 mg/L yeast extract improved both the growth status and tanshinone accumulation. The CT content was increased to a maximum of 2.84 ± 0.33 mg/g DW in the hairy root cultures. Furthermore,

an apparent activation of T-IIA showed a sustained promotion to 2.52 ± 0.67 mg/g DW and DT-I to 1.95 ± 0.09 mg/g DW [65]. The T-I content was slightly responsive to elicitation [65]; however, it was drastically increased to approximately 37.14 times the control level [193].

4.2.2. Abiotic Elicitors

Heavy Metals

In abiotic elicitation, chemical or physical stimuli are used to trigger the synthesis of plant metabolites [195].

Various heavy metal elements have been used to induce specialised metabolites in *S. miltiorrhiza* such as lanthanum, cerium, silver, cobalt, and cadmium [173,196–198], as reported in Table S5 (Supplementary Materials). Among them, Ag^+ was the most frequent elicitor. However, it caused a dose-dependent depression of *S. miltiorrhiza* hairy roots and cell growth [173,198]. A *S. miltiorrhiza* cell suspension exposed to Ag^+ 25 μM for 7 days significantly increased the total tanshinone (CT, T-I, and T-IIA) content to 2.04 mg/g, about 10-fold of the control (0.20 mg/g), with a large amount of CT at 1817.5 $\mu\text{g/g}$, about 30-fold of the control (59.9 $\mu\text{g/g}$) [173]. Recently, Yu et al. [199] showed that on day 7 after treatment with Ag^+ 60 μM , CT was 18.07 times higher than the control.

Plant Signal Compounds

Jasmonates are defined as “hormones” because they elicit cellular responses at low concentrations far from their site of synthesis [200,201]. MJ acts as an efficient elicitor of secondary metabolite production throughout the plant kingdom, particularly those involved in developmental processes and defence responses [202]. It has been used by some researchers to increase the in vitro production of diterpenoids in some *Salvia* species. Some studies on the elicitation of hairy roots of *S. miltiorrhiza* by MJ showed that 100 to 150 μM is the ideal concentration to ensure that the production of various tanshinones is at least four times higher than that of the control [203–205]. However, these results were very different from the previous ones that reported that after elicitation with MJ, the CT and T-IIA were 23.8-fold and 6.2-fold higher, respectively, than those of the control [206]. In cell culture, MJ showed only a moderate or insignificant stimulating effect on tanshinone accumulation in normal and transformed *S. miltiorrhiza* cell cultures [181,188]. MJ showed only a moderate stimulating effect on tanshinone accumulation in *S. castanea* Diels f. *tomentosa* Stib. hairy root culture [65]. Kuźma et al. [207] reported that the total diterpenoid content of *S. sclarea* hairy roots in shake flasks elicited with 125 μM MJ for 7 days was twice that of the control roots. In addition, the diterpenoid content of the same treatment, using a bioreactor as a growth system (67.5 mg/g DW), was about 6-fold higher than that of the non-elicited roots and 2.4-fold higher than that of the MJ-treated roots maintained in shake flasks (28.16 ± 1.2 mg/g DW). Aethiopinone was found to be the major diterpenoid synthesised after MJ treatment among all of the diterpenoids present in *S. sclarea* roots. Its content was 40 mg/g DW, 9.1-fold higher than the control and about 60% of the total diterpenoids. Vaccaro et al. [185] found that 100 μM MJ applied for 7 days to the hairy roots of *S. sclarea* seemed to be more effective in inducing the accumulation of aethiopinone (9.72 ± 0.08 mg/g DW), corresponding to a 25-fold increase over the content of the untreated hairy roots (0.38 ± 0.07 mg/g DW).

Few studies on the genus *Salvia* have used SA elicitation to enhance diterpenoid production (Tables S3 and S6a, Supplementary Materials). SA has been reported as an effective elicitor for the hairy roots and cell cultures of *S. miltiorrhiza*. Hao et al. [204] showed that the total tanshinone production after the application of SA to *S. miltiorrhiza* SmGGPPS, the overexpression of hairy roots was only 1.63-fold that of the mimic treatment control. Zhao et al. [173] and Yu et al. [199] treated the *S. miltiorrhiza* non-transformed cell culture with 100 and 200 μM for 7 days, and the content of CT produced was 6.5-fold and 4.39-fold higher than the control, respectively (Table S3, Supplementary Materials). Kračun-Kolarević et al. [208] reported a significant increase in carnosol and carnosic acid in the shoot cultures of *S. officinalis*. The carnosol and carnosic acid contents were higher

(3.8 and 1.4 times, respectively) in the 4-week-old controls than in 1-week-old explants. SA treatment increased the carnosol production from 2 mg/g DW (in 1-week-old control explants) to 14 mg/g DW (in 4-week-old shoots growing at 150 μ M).

Other Elicitors

Table S6b (Supplementary Materials) report the effect of other chemical compounds on diterpenoid production in in vitro *S. miltiorrhiza* hairy root cultures. The application of 100 μ M sodium nitroprusside (SNP), a donor of NO, resulted in a significant increase in the production levels of T-I, CT, DT-I, and T-IIA in the hairy roots by 80, 170, 60, and 180% above the control level, respectively [209]. The application of β -aminobutyric acid as an elicitor at a dose of 2 mM caused a significant increase in the total tanshinone content of 1.09 mg/g DW, about 4.5 times that of the control at 0.24 mg/g DW [190]. The addition of PEG 2% increased the T-I, CT, DT-I, and T-IIA in the hairy roots to 0.9, 2.1, 3.9, and 2.0 mg/L, respectively, compared to the control at 0.7, 1.3, 3.2, and 1.5 mg/L, respectively [209]. Smoke–water (SW) 1:1000 (*v/v*) was found to be very effective in enhancing the T-I accumulation in hairy roots after 3 days of elicitation [210].

Effect of Light Irradiation

The spectral quality, intensity, and duration of light irradiation may affect plant cell and tissue cultures. Few studies have demonstrated the involvement of light irradiation in the accumulation of secondary metabolites in *Salvia* (Table S7, Supplementary Materials). The hairy roots of *S. miltiorrhiza* exposed to UV-B irradiation showed an increasing production of three tanshinones (CT, T-IA, and T-IIA) and reached a peak (0.38 mg/g DW) that was 1.8 times higher than that of the control. Among the three tanshinones, CT was the most abundant and reached the maximum (0.13 mg/g DW), 3.4 times that of the control, after 40 min of UV-B irradiation [211]. Recently, it was shown that different 1- and 3-week LED light spectrum treatments could regulate tanshinones in the hairy roots of *S. miltiorrhiza*. However, blue light decreased the T-IIA content via the downregulation of key enzymes involved in the biosynthesis process [212].

4.2.3. Combination or Synergic Effect of Elicitors

Some studies have shown that the production of secondary metabolites in *S. miltiorrhiza* hairy root cultures can be enhanced or potentiated by incorporating multiple elicitor treatments over the culture period rather than a single treatment [204,211,213,214]. Many studies have investigated the combination of yeast extract with other elicitors (Tables S8 and S9, Supplementary Materials) in *Salvia* hairy root and cell cultures. The Ti-transformed *S. miltiorrhiza* cell cultures treated with the combination of yeast extract and SA for 15 days increased the total tanshinone production (CT and T-IIA) to the highest levels, with 15.07 mg/L. The yeast extract treatment had 12.23 mg/L and the control and SA treatments had barely detectable amounts of tanshinone [186]. A few years earlier, Chen and Chen [188] showed that in cell suspension, 200 μ M SA added one day before the yeast elicitor increased the CT formation from 13.8 mg/L to 18.9 mg/L (an increase of 37%). Combinations of two elicitors (YE + Ag⁺ or YE + Cd²⁺) and three elicitors (YE + Ag⁺ + Cd²⁺) in *S. miltiorrhiza* cell culture increased the tanshinone content by about 20% and 40%, respectively, compared to a single elicitor [173].

Sorbitol, widely used as an osmotic in plant cell and tissue cultures, has a slight enhancing effect on root growth [85] and can have a positive effect on the production of secondary metabolites [215]. The combination of yeast extract with sorbitol (100 mg/L + 50 g/L, respectively) added on day 21 and maintained for 9 days in suspension in a hairy root culture of *S. miltiorrhiza* decreased the biomass production, and more significantly, increased tanshinone (CT, T-I, and T-IIA) production (16.3 versus 1.77 mg/L) compared to the control [189]. One year later, Wu et al. [191] reported that the combination of yeast extract 25 mg/L with sorbitol +50 g/L increased the biomass production and more drastically increased the tanshinone (CT, T-I, and T-IIA) production by 7-fold (from 0.2 to

1.6 mg/g DW) and the volumetric yield by 13-fold (from 1.95 to 27.4 mg/L) compared to the batch control culture.

Good tanshinone production was obtained by the combination of yeast extract pre-treated for 3 days with β -aminobutyric acid, reaching the amount of 20.1 mg/L [190]. Treatment for 2 days with Ag^+ with CT dramatically increased production [216]. The combination of MJ with a pre-treatment of 40 min with UV-B on hairy roots of *S. miltiorrhiza* significantly increased the total tanshinone content (CT, T-I, and T-IIA) to 28.21 mg/L, 4.9 times higher than that of the control [211].

4.2.4. Elicitation and Nutrient or Medium Feeding or Renewal

Precursor feeding is an obvious and popular approach to increasing the secondary metabolite production in plant cell cultures. Attempts have been made to induce or increase the production of plant secondary metabolites by feeding precursors [149]. In principle, elicitation has been used to induce or stimulate the secondary metabolism of plant cells, but it is not favourable for cell growth (primary metabolism). However, suppressed biomass growth has been observed in hairy root cultures treated with abiotic and biotic elicitors, especially at relatively high doses. This unfavourable effect could be reduced or eliminated during the elicitation process by nutrient or medium supplementation or renewal, thus improving the secondary metabolite production more effectively [192,217–219]. Treatment with 15 μM Ag^+ caused a depression of *S. miltiorrhiza* hairy root (9.6 g DW/L versus 12.1 g DW/L of the control). Supplementation of sucrose and fresh medium prior to the addition of Ag^+ to the culture medium maintained the hairy root growth at 18.9 g DW/L and 22.3 g DW/L versus 12.1 g DW/L of the control, respectively, and the total tanshinone production at 36.6 mg/L and 55.7 mg/L versus 24.4 mg/L and 7.3 mg/L of the Ag^+ treatment and control, respectively [198]. It was found that sorbitol was mainly a strong stimulator of tanshinone production and had beneficial effects on the hairy root growth of *S. miltiorrhiza* [189,191]. To maintain secondary metabolite production throughout the extended fed-batch process without affecting the biomass, both frequent elicitor challenge (sorbitol + YE) and sufficient nutrient supply to the hairy roots of *S. miltiorrhiza* (from 10 mL every 5 days) were essential. Of note was the extremely high tanshinone content achieved in the OS + YE treated culture of 18.1 mg/g DW (equivalent to 1.8% *w/w* or 143.9 mg/L TT yield). This was approximately 11.5 times higher than in the single treatment process (1.57 mg/g DW) and almost 100 times higher than in the absence of any elicitor treatment (0.2 mg/g DW). Moreover, the root fragments released from the roots in the OS + YE treated culture had an even higher tanshinone content (110 mg/g DW or 11% *w/w*) [191].

4.3. Production with In Situ Adsorption

Some secondary metabolites are hydrophobic and are stored intracellularly either in the cytosol or in the cell vacuole. They are minimally secreted into the culture medium and may appear in the culture broth as a result of cell lysis [155,220]. The production of target metabolites can be limited by both the feedback inhibition of accumulated synthesised metabolites and their degraded compounds [221]. However, some secondary metabolites produced in the cell may be released into the culture medium. Miyasaka et al. [222] found that in immobilised cultured cells of *S. miltiorrhiza*, much of the cryptotanshinone was released into the medium, whereas most of the ferruginol was retained in the cells. The introduction of an in situ product removal mechanism such as a solid adsorbent into the culture medium can often effectively induce product release from plant cells and increase productivity [223].

For the enhancement of metabolite production in plant cell culture, different adsorbents for the solid/liquid two-phase system have been used. The addition of 2 g of X-5 resin in a nylon bag to each 200 mL flask recovered a large portion of tanshinones (CT, T-I, and T-IIA) from the roots (70–94%). A significant increase in tanshinone yield was also obtained by combining the macroporous polystyrene resin (X-5 adsorption) with double YE elicitation (added to the culture at 100 mg/L on days 30 and 40). To prolong biomass

growth and secondary metabolite production, medium renewal, YE elicitation, and resin replacement were performed every 10 days for a total of three times (days 30, 40, and 50). Interestingly, the root biomass was increased to 30 g/L DW (versus 8–10 g/L DW in batch mode) and the volumetric tanshinone yield to 87 mg/L (about a 15-fold increase), with 76% adsorbed on the resin. The volumetric productivity of total tanshinone reached 1.46 mg/L per day, more than seven times that of the batch culture [192].

4.4. Gene Overexpression

The overexpression of *SmMDS* in transgenic *S. miltiorrhiza* hairy roots significantly increased the tanshinone yield compared to the control. The total tanshinone content increased in the *SmMDS*-overexpressing lines after elicitor treatment [224]. The overexpression of *SmWRKY1* in *S. miltiorrhiza* significantly increased the transcripts of genes coding for enzymes in the MEP pathway, especially 1-deoxy-D-xylulose-5-phosphate synthase (*SmDXS*) and 1-deoxy-D-xylulose-5-phosphate reductoisomerase (*SmDXR*), resulting in a more than 5-fold increase in tanshinone production in the transgenic lines (up to 13.7 mg/g DW) compared to the control lines [225]. Overexpression of *SmMYB9b* in the hairy roots of *S. miltiorrhiza* increased the tanshinone concentration to 2.16 ± 0.39 mg/g DW, a 2.2-fold improvement over the control [226]. The overexpression of *SmGGPPS* and *SmDXSII* in the hairy roots produced higher levels of tanshinone than the control and single-gene transformed lines; tanshinone production in the double-gene transformed line GDII10 reached 12.93 mg/g DW [227]. Overexpression of *SmAOC* in the hairy roots of *S. miltiorrhiza* significantly increased the yield of T-IIA [228]. Overexpression of the ethylene response factor *SmERF6* in the hairy roots of *S. miltiorrhiza* increased their tanshinone accumulation [229], while the overexpression of the heterologous TFs *AtWRKY18*, *AtWRKY40*, and *AtMYC2* in the hairy roots of *S. sclarea* resulted in the expression of several genes encoding enzymes of the MEP-dependent pathway, in particular *DXS*, *DXR*, *GGPPS*, and *CPPS*. The final yield of aethiopinone was significantly increased by a factor of four in HR lines overexpressing *AtWRKY40* TF [230]. The engineering of the plastidial 2-C-methyl-D-erythritol 4-phosphate-derived isoprenoid pathway in *S. sclarea* hairy roots by the ectopic expression and plastid targeting of cyanobacterial genes encoding the 1-deoxy-D-xylulose 5-phosphate synthase or 1-deoxy-D-xylulose-5-phosphate reductoisomerase gene, the first two enzymatic steps of the plastidial MEP pathway and the plastid-targeted expression of these proteins, significantly increased the yield of aethiopinone by 3-fold and about 6-fold, respectively. The accumulation of other abietane-type diterpenoids (ferruginol, salvipisone, and carnosic acid) was also increased [231].

5. Conclusions

Salvia species are an important source of secondary metabolites and most of these species have been studied in vitro by tissue and cell culture for the controlled production of bioactive diterpenoids. Several techniques have been used to increase the production of cell lines and to optimise the growth medium by using specific precursors or elicitors. This review analysed the biotechnological approaches applied to the in vitro cultures of *Salvia* cells and tissues to enhance the production and accumulation of bioactive diterpenoids and summarised their biological activities. The study showed that culture type, growth medium composition, nutrient supply, elicitors, and in situ uptake can influence both the growth biomass and production of bioactive diterpenoids. All classes of elicitors had a stimulating effect on diterpenoid production in the in vitro cultures of *Salvia* species. *S. miltiorrhiza* was the most studied species for in vitro diterpenoid production, and cryptotanshinone was one of the most stimulated diterpenoids by elicitation. The effects of different elicitors on diterpenoid production in plant tissues and cell cultures were dependent on their concentration, time, and duration of application as well as on the specificity of the secondary metabolites. However, the biomass of *Salvia* cell cultures was more negatively affected by the elicitors compared to tissues. The combination of the overexpression of key genes in a suitable tissue, growing on an ideal medium treated with the best elicitor, with nutrient

feeding and in situ uptake could be a strategy for the production of diterpenoids in *Salvia* species. In conclusion, the application of biotechnological strategies to in vitro cultures of *Salvia* species represents a promising approach for the controlled production of bioactive diterpenoids. These techniques have the potential to revolutionise the pharmaceutical and nutraceutical industries by providing a sustainable and reliable source of valuable medicinal compounds with further research and development including system scale-up. However, further studies are needed to overcome existing challenges and to fully exploit the benefits of these innovative approaches.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/agronomy14040835/s1>, Table S1. Effect of microorganism interaction and their extracts on diterpenoid production in in vitro *Salvia* hairy root cultures; Table S2. Effect of methyl jasmonate and coronatine on diterpenoid production in in vitro *Salvia* hairy root cultures; Table S3. Effect of elicitors on diterpenoid production in in vitro *Salvia* cell cultures; Table S4. Effect of yeast extract on diterpenoid production in in vitro *Salvia* hairy root cultures; Table S5. Effect of Ag⁺ on diterpenoid production in in vitro *Salvia* hairy root cultures; Table S6a. Effect of other chemical compounds on diterpenoid production in in vitro *Salvia* hairy root cultures; Table S6b. Effect of other elicitors on diterpenoid production in in vitro *Salvia* hairy root cultures; Table S7. Effect of physical elicitors on diterpenoid production in in vitro *Salvia* hairy root cultures; Table S8. Effect of the combination of elicitors on diterpenoid production in in vitro *Salvia* hairy root cultures; Table S9. Effect of the combination of elicitors on diterpenoid production in in vitro *Salvia* cell cultures.

Author Contributions: Conceptualisation, R.A.D.K., P.D. and V.I.; Writing—original draft preparation, R.A.D.K., P.D., A.C., V.I., B.R. and V.P.; Writing—review and editing, V.I. and A.B.; Supervision, A.B.; Project administration, A.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflicts of interest.

References

- Will, M.; Claßen-Bockhoff, R. Time to split *Salvia* s.l. (Lamiaceae)—New insights from Old World *Salvia* phylogeny. *Mol. Phylog. Evol.* **2017**, *109*, 33–58. [CrossRef]
- Hu, G.-X.; Takano, A.; Drew, B.T.; Liu, E.-D.; Soltis, D.E.; Soltis, P.S.; Peng, H.; Xiang, C.-L. Phylogeny and staminal evolution of *Salvia* (Lamiaceae, Nepetoideae) in East Asia. *Ann. Bot.* **2018**, *122*, 649–668. [CrossRef]
- Jenks, A.A.; Walker, J.B.; Kim, S.C. Phylogeny of New World *Salvia* subgenus *Calosiphace* (Lamiaceae) based on cpDNA (psbA-trnH) and nrDNA (ITS) sequence data. *J. Plant Res.* **2013**, *126*, 483–496. [CrossRef]
- Drew, B.T.; González-Gallegos, J.G.; Xiang, C.-L.; Kriebel, R.; Drummond, C.P.; Walker, J.B.; Sytsma, K.J. *Salvia* united: The greatest good for the greatest number. *Taxon* **2017**, *66*, 133–145. [CrossRef]
- Hardman, R. *SAGE—The Genus Salvia*; Kintzios, S.E., Ed.; OPA (Overseas Publishers Association) N.V.; Harwood Academic Publishers, Part of the Gordon and Breach Publishing Group: New York, NY, USA, 2000.
- Hamidpour, M.; Hamidpour, R.; Hamidpour, S.; Shahlari, M. Chemistry, pharmacology, and medicinal property of sage (*Salvia*) to prevent and cure illnesses such as obesity, diabetes, depression, dementia, lupus, autism, heart disease, and cancer. *J. Tradit. Complement. Med.* **2014**, *4*, 82–88. [CrossRef]
- Bisio, A.; Pedrelli, F.; D’Ambola, M.; Labanca, F.; Schito, A.M.; Govaerts, R.; De Tommasi, N.; Milella, L. Quinone diterpenes from *Salvia* species: Chemistry, botany, and biological activity. *Phytochem. Rev.* **2019**, *18*, 665–842. [CrossRef]
- Wu, Y.-B.; Ni, Z.-Y.; Shi, Q.-W.; Dong, M.; Kiyota, H.; Gu, Y.-C.; Cong, B. Constituents from *Salvia* species and their biological activities. *Chem. Rev.* **2012**, *112*, 5967–6026. [CrossRef]
- Xu, J.; Wei, K.; Zhang, G.; Lei, L.; Yang, D.; Wang, W.; Han, Q.; Xia, Y.; Bi, Y.; Yang, M.; et al. Ethnopharmacology, phytochemistry, and pharmacology of Chinese *Salvia* species: A review. *J. Ethnopharmacol.* **2018**, *225*, 18–30. [CrossRef]
- Iobbi, V.; Brun, P.; Bernabé, G.; Dougué Kentsop, R.A.; Donadio, G.; Ruffoni, B.; Fossa, P.; Bisio, A.; De Tommasi, N. Labdane diterpenoids from *Salvia tingitana* Etl. Synergize with clindamycin against methicillin-resistant *Staphylococcus aureus*. *Molecules* **2021**, *26*, 6681. [CrossRef]
- Lubbe, A.; Verpoorte, R. Cultivation of medicinal and aromatic plants for specialty industrial materials. *Ind. Crop. Prod.* **2011**, *34*, 785–801. [CrossRef]

12. FDA. Title 21—Food and Drugs. Chapter I—Food and Drug Administration. Department of Health and Human Services. Subchapter B—Food for Human Consumption (Continued). PART 182—Substances Generally Recognized as Safe. Subpart A—General Provisions. Sec. 182.10 Spices and Other Natural Seasonings and Flavorings; FDA: Washington, DC, USA, 2023.
13. Bisio, A.; Schito, A.M.; Pedrelli, F.; Danton, O.; Reinhardt, J.K.; Poli, G.; Tuccinardi, T.; Bürgi, T.; De Riccardis, F.; Giacomini, M.; et al. Antibacterial and ATP synthesis modulating compounds from *Salvia tingitana*. *J. Nat. Prod.* **2020**, *83*, 1027–1042. [\[CrossRef\]](#)
14. Topçu, G. Bioactive triterpenoids from *Salvia* species. *J. Nat. Prod.* **2006**, *69*, 482–487. [\[CrossRef\]](#)
15. Topçu, G.; Ozturk, M.; Kusman, T.; Barla Demirköz, A.A.; Kolak, U.; Ulubelen, A. Terpenoids, essential oil composition, fatty acid profile, and biological activities of Anatolian *Salvia fruticosa* Mill. *Turk. J. Chem.* **2013**, *37*, 619–632. [\[CrossRef\]](#)
16. Rodríguez-Hahn, L.; Esquivel, B.; Cárdenas, J.; Ramamoorthy, T.P. The distribution of diterpenoids in *Salvia*. In *Advances in Labiatae Science*; Harley, R.M., Reynolds, T., Eds.; The Royal Botanic Gardens: Richmond, UK, 1992.
17. Lu, Y.R.; Foo, L.Y. Polyphenolics of *Salvia*—A review. *Phytochemistry* **2002**, *59*, 117–140. [\[CrossRef\]](#)
18. Li, M.-H.; Chen, J.-M.; Peng, Y.; Xiao, P.-G. Distribution of phenolic acids in Chinese *Salvia* plants. *WST* **2008**, *10*, 46–52.
19. Valant-Vetschera, K.M.; Roitman, J.N.; Wollenweber, E. Chemodiversity of exudate flavonoids in some members of the Lamiaceae. *Biochem. Syst. Ecol.* **2003**, *31*, 1279–1289. [\[CrossRef\]](#)
20. Tomas-Barberan, F.A.; Wollenweber, E. Flavonoid aglycones from the leaf surfaces of some *Labiatae* species. *Plant Syst. Evol.* **1990**, *173*, 109–118. [\[CrossRef\]](#)
21. Kabouche, A.; Kabouche, Z. Bioactive Diterpenoids of *Salvia* Species. *Stud. Nat. Prod. Chem.* **2008**, *35*, 753–833.
22. Bisio, A.; Fontana, N.; Romussi, G.; De Tommasi, N. Diterpenes and triterpenes from *Salvia aurea* L.—Part-2—Constituents of *Labiatae*. *Pharmazie* **1998**, *53*, 210–211.
23. Rodríguez-Hahn, L.; Esquivel, B.; Sanchez, A.; Sanchez, C.; Cardenas, J.; Ramamoorthy, T. Diterpenos abietánicos de *Salvias* mexicanas. *Rev. Latinoamer. Quím.* **1989**, *20*, 105–110.
24. Hu, J.; Wang, F.; Liang, F.; Wu, Z.; Jiang, R.; Li, J.; Chen, J.; Qiu, S.; Wang, J.; Zhang, Y.; et al. Identification of abietane-type diterpenoids and phenolic acids biosynthesis genes in *Salvia apiana* Jepson through full-length transcriptomic and metabolomic profiling. *Front. Plant Sci.* **2022**, *13*, 919025. [\[CrossRef\]](#)
25. Rodríguez-Hahn, L.; Esquivel, B.; Cárdenas, J. Clerodane diterpenes in *Labiatae*. In *Progress in the Chemistry of Organic Natural Products*; Herz, W., Kirby, G.W., Moore, R.E., Steglich, W., Tamm, C., Eds.; Springer: Vienna, Austria, 1994; Volume 63, pp. 107–196.
26. Bisio, A.; Damonte, G.; Fraternali, D.; Giacomelli, E.; Sails, A.; Romussi, G.; Cafaggi, S.; Ricci, D.; De Tommasi, N. Phytotoxic clerodane diterpenes from *Salvia miniata* Fernald (Lamiaceae). *Phytochemistry* **2011**, *72*, 265–275. [\[CrossRef\]](#)
27. Bisio, A.; Mieri, M.D.; Milella, L.; Schito, A.M.; Parricchi, A.; Russo, D.; Alfei, S.; Lapillo, M.; Tuccinardi, T.; Hamburger, M. Antibacterial and hypoglycemic diterpenoids from *Salvia chamaedryoides*. *J. Nat. Prod.* **2017**, *80*, 503–514. [\[CrossRef\]](#)
28. Bisio, A.; Fontana, N.; Romussi, G.; Ciarallo, G.; De Tommasi, N.; Pizza, C.; Mugnoli, A. Clerodane diterpenoids from *Salvia blepharophylla*. *Phytochemistry* **1999**, *52*, 1535–1540. [\[CrossRef\]](#)
29. Capasso, R.; Izzo, A.; Capasso, F.; Romussi, G.; Bisio, A.; Mascolo, N. A diterpenoid from *Salvia cinnabarina* inhibits mouse intestinal motility in vivo. *Planta Medica* **2004**, *70*, 375–377. [\[CrossRef\]](#)
30. Bisio, A.; Romussi, G.; Russo, E.; Cafaggi, S.; Schito, A.M.; Repetto, B.; De Tommasi, N. Antimicrobial activity of the ornamental species *Salvia corrugata*, a potential new crop for extractive purposes. *J. Agric. Food Chem.* **2008**, *56*, 10468–10472. [\[CrossRef\]](#)
31. Giacomelli, E.; Bertrand, S.; Nievergelt, A.; Zwick, V.; Simoes-Pires, C.; Marcourt, L.; Rivara-Minten, E.; Cuendet, M.; Bisio, A.; Wolfender, J.-L. Cancer chemopreventive diterpenes from *Salvia corrugata*. *Phytochemistry* **2013**, *96*, 257–264. [\[CrossRef\]](#)
32. Simmons, E.M.; Sarpong, R. Structure, biosynthetic relationships and chemical synthesis of the icetexane diterpenoids. *Nat. Prod. Rep.* **2009**, *26*, 1195–1217. [\[CrossRef\]](#)
33. Marchev, A.; Haas, C.; Schulz, S.; Georgiev, V.; Steingroewer, J.; Bley, T.; Pavlov, A. Sage in vitro cultures: A promising tool for the production of bioactive terpenes and phenolic substances. *Biotechnol. Lett.* **2014**, *36*, 211–221. [\[CrossRef\]](#)
34. Karalija, E.; Dahija, S.; Tarkowski, P.; Zeljković, S. Influence of climate-related environmental stresses on economically important essential oils of Mediterranean *Salvia* spp. *Front. Plant Sci.* **2022**, *13*, 864807. [\[CrossRef\]](#)
35. Mohaddab, M.; El Goumi, Y.; Gallo, M.; Montesano, D.; Zengin, G.; Bouyahya, A.; Fakiri, M. Biotechnology and in vitro culture as an alternative system for secondary metabolite production. *Molecules* **2022**, *27*, 8093. [\[CrossRef\]](#)
36. Ozyigit, I.; Dogan, I.; Hocaoglu-Ozyigit, A.; Yalcin, B.; Erdogan, A.; Yalcin, I.E.; Cabi, E.; Kaya, Y. Production of secondary metabolites using tissue culture-based biotechnological applications. *Front. Plant Sci.* **2023**, *14*, 1132555. [\[CrossRef\]](#)
37. Grzegorzczak, I.; Bilichowski, I.; Mikiciuk-Olasik, E.; Wysokinska, H. In vitro cultures of *Salvia officinalis* L. as a source of antioxidant compounds. *Acta Soc. Bot. Pol.* **2005**, *74*, 17–21. [\[CrossRef\]](#)
38. Bisio, A.; Fraternali, D.; Schito, A.M.; Parricchi, A.; Dal Piaz, F.; Ricci, D.; Giacomini, M.; Ruffoni, B.; De Tommasi, N. Establishment and analysis of in vitro biomass from *Salvia corrugata* Vahl. and evaluation of antimicrobial activity. *Phytochemistry* **2016**, *122*, 276–285. [\[CrossRef\]](#)
39. Ruffoni, B.; Raffi, D.; Rizzo, A.M.; Oleszek, W.; Giardi, M.T.; Bertoli, A.; Pistelli, L. Establishment of in vitro *Salvia* cell biomass for the controlled production of antioxidant metabolites. *Acta Hort.* **2010**, *829*, 423–427. [\[CrossRef\]](#)
40. Martini, A.N.; Vlachou, G.; Papafotiou, M. Effect of explant origin and medium plant growth regulators on in vitro shoot proliferation and rooting of *Salvia tomentosa*, a native sage of the Northeastern Mediterranean basin. *Agronomy* **2022**, *12*, 1889. [\[CrossRef\]](#)
41. Georgiev, V.; Pavlov, A. *Salvia Biotechnology*; Springer: New York, NY, USA, 2017; pp. 1–439.

42. Kintzios, S.; Nikolaou, A.; Skoula, M. Somatic embryogenesis and in vitro rosmarinic acid accumulation in *Salvia officinalis* and *S. fruticosa* leaf callus cultures. *Plant Cell Rep.* **1999**, *18*, 462–466. [\[CrossRef\]](#)
43. Bassolino, L.; Giacomelli, E.; Giovanelli, S.; Pistelli, L.; Cassetti, A.; Damonte, G.; Bisio, A.; Ruffoni, B. Tissue culture and aromatic profile in *Salvia dolomitica* Codd. *Plant Cell Tissue Organ Cult.* **2014**, *121*, 83–95. [\[CrossRef\]](#)
44. Fraternali, D.; Bisio, A.; Ricci, D. *Salvia x jamensis* J. Compton: In vitro regeneration of shoots through TDZ and BA. *Plant Biosyst.* **2013**, *147*, 713–716. [\[CrossRef\]](#)
45. Dougué Kentsop, R.A. *Biotechnology Applied to Aromatic Plants for the Controlled Production of Bioactive Compounds*; University of Genoa: Genoa, Italy, 2020.
46. Hanson, J.R. Diterpenoids. *Nat. Prod. Rep.* **1986**, *3*, 307–322. [\[CrossRef\]](#) [\[PubMed\]](#)
47. Hanson, J.R. Diterpenoids. *Nat. Prod. Rep.* **1987**, *4*, 399–413. [\[CrossRef\]](#)
48. Hanson, J.R. Diterpenoids. *Nat. Prod. Rep.* **1990**, *7*, 149–164. [\[CrossRef\]](#)
49. Hanson, J.R. Diterpenoids. *Nat. Prod. Rep.* **1992**, *9*, 1–16. [\[CrossRef\]](#)
50. Hanson, J.R. Diterpenoids. *Nat. Prod. Rep.* **1994**, *11*, 265–277. [\[CrossRef\]](#)
51. Hanson, J.R. Diterpenoids. *Nat. Prod. Rep.* **1996**, *13*, 59–71. [\[CrossRef\]](#)
52. Hanson, J.R. Diterpenoids. *Nat. Prod. Rep.* **2002**, *19*, 125–132. [\[CrossRef\]](#)
53. Hanson, J.R. Diterpenoids. *Nat. Prod. Rep.* **2003**, *20*, 70–78. [\[CrossRef\]](#)
54. Hanson, J.R. Diterpenoids. *Nat. Prod. Rep.* **2004**, *21*, 785–793. [\[CrossRef\]](#)
55. Hanson, J.R. Diterpenoids. *Nat. Prod. Rep.* **2005**, *22*, 594–602. [\[CrossRef\]](#) [\[PubMed\]](#)
56. Hanson, J.R. Diterpenoids. *Nat. Prod. Rep.* **2006**, *23*, 875–885. [\[CrossRef\]](#) [\[PubMed\]](#)
57. Hanson, J.R. Diterpenoids. *Nat. Prod. Rep.* **2009**, *26*, 1156–1171. [\[CrossRef\]](#)
58. Hanson, J.R. Diterpenoids of terrestrial origin. *Nat. Prod. Rep.* **2011**, *28*, 1755–1772. [\[CrossRef\]](#)
59. Hanson, J.R. Diterpenoids of terrestrial origin. *Nat. Prod. Rep.* **2015**, *32*, 1654–1663. [\[CrossRef\]](#)
60. Hanson, J.R. Diterpenoids of terrestrial origin. *Nat. Prod. Rep.* **2016**, *33*, 1227–1238. [\[CrossRef\]](#)
61. Hanson, J.R. Diterpenoids of terrestrial origin. *Nat. Prod. Rep.* **2017**, *34*, 1233–1243. [\[CrossRef\]](#)
62. Yang, M.H.; Blunden, G.; Xu, Y.X.; Nagy, G.; Mathe, I. Diterpenoids from *Salvia* species. *Pharm. Pharmacol. Commun.* **1996**, *2*, 69–71.
63. Zhi, B.H.; Alfermann, A. Diterpenoid production in hairy root cultures of *Salvia miltiorrhiza*. *Phytochemistry* **1993**, *32*, 699–703. [\[CrossRef\]](#)
64. Chen, H.; Yuan, J.-P.; Chen, F.; Zhang, Y.-L.; Song, J.-Y. Tanshinone production in Ti-transformed *Salvia miltiorrhiza* cell suspension cultures. *J. Biotechnol.* **1997**, *58*, 147–156. [\[CrossRef\]](#)
65. Li, B.; Wang, B.; Li, H.; Peng, L.; Ru, M.; Liang, Z.; Yan, X.; Zhu, Y. Establishment of *Salvia castanea* Diels f. *tomentosa* Stib. hairy root cultures and the promotion of tanshinone accumulation and gene expression with Ag(+), methyl jasmonate, and yeast extract elicitation. *Protoplasma* **2016**, *253*, 87–100. [\[CrossRef\]](#)
66. Vaccaro, M.; Malafronte, N.; Alfieri, M.; De Tommasi, N.; Leone, A. Enhanced biosynthesis of bioactive abietane diterpenes by overexpressing AtDXS or AtDXR genes in *Salvia sclarea* hairy roots. *Plant Cell Tissue Organ Cult.* **2014**, *119*, 65–77. [\[CrossRef\]](#)
67. Santos-Gomes, P.C.; Seabra, R.M.; Andrade, P.B.; Fernandes-Ferreira, M. Determination of phenolic antioxidant compounds produced by calli and cell suspensions of sage (*Salvia officinalis* L.). *J. Plant Physiol.* **2003**, *160*, 1025–1032. [\[CrossRef\]](#)
68. Kuźma, Ł.; Kaiser, M.; Wysokińska, H. The production and antiprotozoal activity of abietane diterpenes in *Salvia austriaca* hairy roots grown in shake flasks and bioreactor. *Prep. Biochem. Biotechnol.* **2017**, *47*, 58–66. [\[CrossRef\]](#)
69. Li, M.H.; Peng, Y.; Xiao, P.G. Distribution of tanshinones in the genus *Salvia* (family Lamiaceae) from China and its systematic significance. *J. Syst. Evol.* **2010**, *48*, 118–122. [\[CrossRef\]](#)
70. Zhang, Y.; Jiang, P.; Ye, M.; Kim, S.-H.; Jiang, C.; Lü, J. Tanshinones: Sources, Pharmacokinetics and Anti-Cancer Activities. *Int. J. Mol. Sci.* **2012**, *13*, 13621–13666. [\[CrossRef\]](#)
71. Li, M.-H.; Li, Q.-Q.; Liu, Y.-Z.; Cui, Z.-H.; Zhang, N.; Huang, L.-Q.; Xiao, P.-G. Pharmacophylogenetic study on plants of genus *Salvia* L. from China. *Chin. Herb. Med.* **2013**, *5*, 164–181.
72. Zhou, L.; Zuo, Z.; Chow, M.S.S. Danshen: An overview of its chemistry, pharmacology, pharmacokinetics, and clinical use. *J. Clin. Pharmacol.* **2005**, *45*, 1345–1359. [\[CrossRef\]](#)
73. Li, H.-B.; Chen, F. Preparative isolation and purification of six diterpenoids from the Chinese medicinal plant *Salvia miltiorrhiza* by high-speed counter-current chromatography. *J. Chromatogr. A* **2001**, *925*, 109–114. [\[CrossRef\]](#)
74. Li, M.-H.; Chen, J.-M.; Peng, Y.; Wu, Q.; Xiao, P.-G. Investigation of Danshen and related medicinal plants in China. *J. Ethnopharmacol.* **2008**, *120*, 419–426. [\[CrossRef\]](#)
75. Guo, J.; Zhou, Y.J.; Hillwig, M.L.; Shen, Y.; Yang, L.; Wang, Y.; Zhang, X.; Liu, W.; Peters, R.J.; Chen, X. CYP76AH1 catalyzes turnover of miltiradiene in tanshinones biosynthesis and enables heterologous production of ferruginol in yeasts. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 12108–12113. [\[CrossRef\]](#)
76. Gao, W.; Hillwig, M.L.; Huang, L.; Cui, G.; Wang, X.; Kong, J.; Yang, B.; Peters, R.J. A functional genomics approach to tanshinone biosynthesis provides stereochemical insights. *Org. Lett.* **2009**, *11*, 5170–5173. [\[CrossRef\]](#)
77. Ikeshiro, Y.; Mase, I.; Tomita, Y. Abietane type diterpenoids from *Salvia miltiorrhiza*. *Phytochemistry* **1989**, *28*, 3139–3141. [\[CrossRef\]](#)
78. Lee, A.R.; Wu, W.L.; Chang, W.L.; Lin, H.C.; King, M.L. Isolation and bioactivity of new tanshinones. *J. Nat. Prod.* **1987**, *50*, 157–160. [\[CrossRef\]](#)

79. Wang, X.; Morris-Natschke, S.L.; Lee, K.-H. New developments in the chemistry and biology of the bioactive constituents of tanshen. *Med. Res. Rev.* **2007**, *27*, 133–148. [\[CrossRef\]](#)
80. Bo, L.; Fan-Di, N.; Zhong-Wen, L.; Hong-Jie, Z.; De-Zu, W.; Han-Dong, S. Diterpenoids from the roots of *Salvia przewalskii*. *Phytochemistry* **1991**, *30*, 3815–3817. [\[CrossRef\]](#)
81. Liu, L.; Yang, D.; Xing, B.; Zhang, H.; Liang, Z. *Salvia castanea* hairy roots are more tolerant to phosphate deficiency than *Salvia miltiorrhiza* hairy roots based on the secondary metabolism and antioxidant defenses. *Molecules* **2018**, *23*, 1132. [\[CrossRef\]](#)
82. Skała, E.; Mielicki, W.; Wysokińska, H. Tanshinones in culture of *Salvia przewalskii* Maxim in vitro. *Acta Biol. Cracoviensia Ser. Bot.* **2014**, *56*, 104–110. [\[CrossRef\]](#)
83. Skała, E.; Wysokińska, H. Tanshinone production in roots of micropropagated *Salvia przewalskii* Maxim. *Z. Naturforschung C* **2005**, *60*, 583–586. [\[CrossRef\]](#)
84. Sairafianpour, M.; Christensen, J.; Stærk, D.; Budnik, B.A.; Kharazmi, A.; Bagherzadeh, K.; Jaroszewski, J.W. Leishmanicidal, antiplasmodial, and cytotoxic activity of novel diterpenoid 1,2-quinones from *Perovskia abrotanoides*: New source of tanshinones. *J. Nat. Prod.* **2001**, *64*, 1398–1403. [\[CrossRef\]](#)
85. Zaker, A.; Sykora, C.; Gössnitzer, F.; Abrishamchi, P.; Asili, J.; Mousavi, S.H.; Wawrosch, C. Effects of some elicitors on tanshinone production in adventitious root cultures of *Perovskia abrotanoides* Karel. *Ind. Crop. Prod.* **2015**, *67*, 97–102. [\[CrossRef\]](#)
86. Jiang, Z.; Gao, W.; Huang, L. Tanshinones, critical pharmacological components in *Salvia miltiorrhiza*. *Front. Pharmacol.* **2019**, *10*, 202. [\[CrossRef\]](#)
87. Rodríguez, B.; Fernández-Gadea, F.; Savona, G. A rearranged abietane diterpenoid from the root of *Salvia aethiopis*. *Phytochemistry* **1984**, *23*, 1805–1806. [\[CrossRef\]](#)
88. Ulubelen, A.; Tan, N. Terpenoids from *Salvia recognita* and *Salvia aethiopis*. *Sci. Pharm.* **1999**, *67*, 83–88.
89. Boya, M.T.; Valverde, S. An orthoquinone isolated from *Salvia aethiopis*. *Phytochemistry* **1981**, *20*, 1367–1368. [\[CrossRef\]](#)
90. Michavila, A.; María, C.; Rodríguez, B. 20-Nor-abietane and rearranged abietane diterpenoids from the root of *Salvia argentea*. *Phytochemistry* **1986**, *25*, 1935–1937. [\[CrossRef\]](#)
91. Ulubelen, A.; Topcu, G.; Tan, N. Rearranged abietane diterpenes from *Salvia candidissima*. *Phytochemistry* **1992**, *31*, 3637–3638. [\[CrossRef\]](#)
92. Goren, A.C.; Topcu, G.; Oksuz, S.; Kokdil, G.; Voelter, W.; Ulubelen, A. Diterpenoids from *Salvia ceratophylla*. *Nat. Prod. Lett.* **2002**, *16*, 47–52. [\[CrossRef\]](#)
93. Gökdil, G.; Topcu, G.; Sönmez, U.; Ulubelen, A. Terpenoids and flavonoids from *Salvia cyanescens*. *Phytochemistry* **1997**, *46*, 799–800. [\[CrossRef\]](#)
94. Ulubelen, A.; Sönmez, U.; Topcu, G. Diterpenoids from the roots of *Salvia sclarea*. *Phytochemistry* **1997**, *44*, 1297–1299. [\[CrossRef\]](#)
95. Kuźma, Ł.; Skrzypek, Z.; Wysokińska, H. Diterpenoids and triterpenoids in hairy roots of *Salvia sclarea*. *Plant Cell Tissue Organ Cult.* **2006**, *84*, 171–179. [\[CrossRef\]](#)
96. Kuzma, Ł.; Bruchajze, E.; Wysokińska, H. Diterpenoid production in hairy root culture of *Salvia sclarea* L. *Z. Naturforsch. Sect. C* **2008**, *63*, 621–624. [\[CrossRef\]](#)
97. Kentsop, R.A.D.; Iobbi, V.; Donadio, G.; Ruffoni, B.; De Tommasi, N.; Bisio, A. Abietane diterpenoids from the hairy roots of *Salvia corrugata*. *Molecules* **2021**, *26*, 5144. [\[CrossRef\]](#)
98. Kuźma, Ł.; Derda, M.; Hadaś, E.; Wysokińska, H. Abietane diterpenoids from *Salvia sclarea* transformed roots as growth inhibitors of pathogenic *Acanthamoeba* spp. *Parasitol. Res.* **2015**, *114*, 323–327. [\[CrossRef\]](#)
99. Ebrahimi, S.N.; Zimmermann, S.; Zaugg, J.; Smiesko, M.; Brun, R.; Hamburger, M. Abietane diterpenoids from *Salvia sahendicab*—Antiprotozoal activity and determination of their absolute configurations. *Planta Medica* **2013**, *29*, 150–156.
100. Hernández-Pérez, M.; Rabanal, R.M.; de la Torre, M.C.; Rodríguez, B. Analgesic, anti-inflammatory, antipyretic and haematological effects of aethiopinone, an o-naphthoquinone diterpenoid from *Salvia aethiopis* roots and two hemisynthetic derivatives. *Planta Medica* **1995**, *61*, 505–509. [\[CrossRef\]](#)
101. Walencka, E.; Rozalska, S.; Wysokińska, H.; Rozalski, M.; Kuzma, Ł.; Rozalska, B. Salvipisone and aethiopinone from *Salvia sclarea* hairy roots modulate staphylococcal antibiotic resistance and express anti-biofilm activity. *Planta Medica* **2007**, *73*, 545–551. [\[CrossRef\]](#)
102. Różalski, M.; Walencka, E.; Różalska, B.; Wysokińska, H. Antimicrobial activity of diterpenoids from hairy roots of *Salvia sclarea* L.: Salvipisone as a potential anti-biofilm agent active against antibiotic resistant Staphylococci. *Phytomedicine* **2007**, *14*, 31–35.
103. Hernandez-Perez, M.; Rabanal, R.; Arias, A.; de La Torre, M.; Rodriguez, B. Aethiopinone, an antibacterial and cytotoxic agent from *Salvia aethiopis* roots. *Pharm. Biol.* **1999**, *37*, 17–21. [\[CrossRef\]](#)
104. Różalski, M.; Kuźma, Ł.; Wysokińska, H.; Krajewska, U. Cytotoxic and proapoptotic activity of diterpenoids from in vitro cultivated *Salvia sclarea* roots. Studies on the leukemia cell lines. *Z. Naturforschung C* **2006**, *61*, 483–488. [\[CrossRef\]](#)
105. Božić, D.; Papaefthimiou, D.; Brückner, K.; De Vos, R.C.H.; Tsoleridis, C.A.; Katsarou, D.; Papanikolaou, A.; Pateraki, I.; Chatzopoulou, F.M.; Dimitriadou, E. Towards elucidating carnosic acid biosynthesis in Lamiaceae: Functional characterization of the three first steps of the pathway in *Salvia fruticosa* and *Rosmarinus officinalis*. *PLoS ONE* **2015**, *10*, e0124106. [\[CrossRef\]](#)
106. Birtić, S.; Dussort, P.; Pierre, F.-X.; Bily, A.C.; Roller, M. Carnosic acid. *Phytochemistry* **2015**, *115*, 9–19. [\[CrossRef\]](#)
107. Lin, K.-I.; Lin, C.-C.; Kuo, S.-M.; Lai, J.-C.; Wang, Y.-Q.; You, H.-L.; Hsu, M.-L.; Chen, C.-H.; Shiu, L.-Y. Carnosic acid impedes cell growth and enhances anticancer effects of carmustine and lomustine in melanoma. *Biosci. Rep.* **2018**, *38*, BSR20180005. [\[CrossRef\]](#)

108. Pavić, V.; Jakovljević, M.; Molnar, M.; Jokić, S. Extraction of carnosic acid and carnosol from sage (*Salvia officinalis* L.) leaves by supercritical fluid extraction and their antioxidant and antibacterial activity. *Plants* **2019**, *8*, 16. [\[CrossRef\]](#)
109. PoECKel, D.; Greiner, C.; Verhoff, M.; Rau, O.; Tausch, L.; Hörnig, C.; Steinhilber, D.; Schubert-Zsilavec, M.; Werz, O. Carnosic acid and carnosol potently inhibit human 5-lipoxygenase and suppress pro-inflammatory responses of stimulated human polymorphonuclear leukocytes. *Biochem. Pharmacol.* **2008**, *76*, 91–97. [\[CrossRef\]](#)
110. Bauer, J.; Kuehn, S.; Rollinger, J.M.; Scherer, O.; Northoff, H.; Stuppner, H.; Werz, O.; Koeberle, A. Carnosol and carnosic acids from *Salvia officinalis* inhibit microsomal prostaglandin E2 synthase-1. *J. Pharmacol. Exp. Ther.* **2012**, *342*, 169–176. [\[CrossRef\]](#)
111. Ninomiya, K.; Matsuda, H.; Shimoda, H.; Nishida, N.; Kasajima, N.; Yoshino, T.; Morikawa, T.; Yoshikawa, M. Carnosic acid, a new class of lipid absorption inhibitor from sage. *Bioorganic Med. Chem. Lett.* **2004**, *14*, 1943–1946. [\[CrossRef\]](#)
112. Zunin, P.; Leardi, R.; Bisio, A.; Boggia, R.; Romussi, G. Oxidative stability of virgin olive oil enriched with carnosic acid. *Food Res. Int.* **2010**, *43*, 1511–1516. [\[CrossRef\]](#)
113. Cortese, K.; Daga, A.; Monticone, M.; Tavella, S.; Stefanelli, A.; Aiello, C.; Bisio, A.; Bellese, G.; Castagnola, P. Carnosic acid induces proteasomal degradation of Cyclin B1, RB and SOX2 along with cell growth arrest and apoptosis in GBM cells. *Phytomedicine* **2016**, *23*, 679–685. [\[CrossRef\]](#)
114. Maione, F.; Cantone, V.; Pace, S.; Chini, M.G.; Bisio, A.; Romussi, G.; Pieretti, S.; Werz, O.; Koeberle, A.; Mascolo, N. Anti-inflammatory and analgesic activity of carnosol and carnosic acid in vivo and in vitro and in silico analysis of their target interactions. *Br. J. Pharmacol.* **2017**, *174*, 1497–1508. [\[CrossRef\]](#)
115. D'Alesio, C.; Bellese, G.; Gagliani, M.C.; Aiello, C.; Grasselli, E.; Marcocci, G.; Bisio, A.; Tavella, S.; Daniele, T.; Cortese, K.; et al. Cooperative antitumor activities of carnosic acid and Trastuzumab in ERBB2+ breast cancer cells. *J. Exp. Clin. Cancer Res.* **2017**, *36*, 154. [\[CrossRef\]](#)
116. Gonzalez, M.A. Aromatic abietane diterpenoids: Their biological activity and synthesis. *Nat. Prod. Rep.* **2015**, *32*, 684–704. [\[CrossRef\]](#)
117. Abreu, M.E.; Müller, M.; Alegre, L.; Munné-Bosch, S. Phenolic diterpene and α -tocopherol contents in leaf extracts of 60 *Salvia* species. *J. Sci. Food Agric.* **2008**, *88*, 2648–2653. [\[CrossRef\]](#)
118. Brückner, K.; Božić, D.; Manzano, D.; Papaefthimiou, D.; Pateraki, I.; Scheler, U.; Ferrer, A.; de Vos, R.C.H.; Kanellis, A.K.; Tissier, A. Characterization of two genes for the biosynthesis of abietane-type diterpenes in rosemary (*Rosmarinus officinalis*) glandular trichomes. *Phytochemistry* **2014**, *101*, 52–64. [\[CrossRef\]](#)
119. Anwar, F.; Qadir, R. Carnosic acid and carnosol. In *A Century of Valuable Plant Bioactives*; Mushtaq, M., Anwar, F., Eds.; Elsevier: Cambridge, MA, USA, 2021; pp. 261–274.
120. Min, F.; Liu, X.; Li, Y.; Dong, M.; Qu, Y.; Liu, W. Carnosic acid suppresses the development of oral squamous cell carcinoma via mitochondrial-mediated apoptosis. *Front. Oncol.* **2021**, *11*, 760861. [\[CrossRef\]](#)
121. Iobbi, V.; Parisi, V.; Bernabè, G.; De Tommasi, N.; Bisio, A.; Brun, P. Anti-biofilm activity of carnosic acid from *Salvia rosmarinus* against methicillin-resistant *Staphylococcus aureus*. *Plants* **2023**, *12*, 3679. [\[CrossRef\]](#)
122. Iobbi, V.; Donadio, G.; Lanteri, A.P.; Maggi, N.; Kirchmair, J.; Parisi, V.; Minuto, G.; Copetta, A.; Giacomini, M.; Bisio, A.; et al. Targeted metabolite profiling of *Salvia rosmarinus* Italian local ecotypes and cultivars and inhibitory activity against *Pectobacterium carotovorum* subsp. *carotovorum*. *Front. Plant Sci.* **2024**, *15*, 1164859. [\[CrossRef\]](#)
123. Jassbi, A.R.; Hadavand Mirzaei, H.; Firuzi, O.; Pirhadi, S.; Asadollahi, M.; Chandran, J.N.; Schneider, B. Cytotoxic abietane-type diterpenoids from roots of *Salvia spinosa* and their in silico pharmacophore modeling. *Nat. Prod. Res.* **2022**, *36*, 3183–3188. [\[CrossRef\]](#)
124. Kuźma, Ł.; Wysokińska, H.; Sikora, J.; Olszewska, P.; Mikiciuk-Olasik, E.; Szymański, P. Taxodione and extracts from *Salvia austriaca* roots as human cholinesterase inhibitors. *Phytother. Res.* **2016**, *30*, 234–242. [\[CrossRef\]](#)
125. Búfalo, J.; Cantrell, C.L.; Jacob, M.R.; Schrader, K.K.; Tekwani, B.L.; Kustova, T.S.; Ali, A.; Boaro, C.S. Antimicrobial and antileishmanial activities of diterpenoids isolated from the roots of *Salvia deserta*. *Planta Medica* **2015**, *82*, 131–137. [\[CrossRef\]](#)
126. Shafaei-Bajestani, N.; Emami, S.A.; Asili, J.; Tayarani-Najaran, Z. Anti-apoptotic effect of taxodione on serum/glucose deprivation-induced PC12 cells death. *Cell. Mol. Neurobiol.* **2014**, *34*, 1103–1109. [\[CrossRef\]](#)
127. Sadowska, B.; Kuźma, Ł.; Micota, B.; Budzyńska, A.; Wysokińska, H.; Kłys, A.; Więckowska-Szakiel, M.; Różalska, B. New biological potential of abietane diterpenoids isolated from *Salvia austriaca* against microbial virulence factors. *Microb. Pathog.* **2016**, *98*, 132–139. [\[CrossRef\]](#)
128. Rodríguez-Hahn, L.; Esquivel, B.; Sánchez, C.; Estebanes, L.; Cárdenas, J.; Soriano-García, M.; Toscano, R.; Ramamoorthy, T. Abietane type diterpenoids from *Salvia fruticulosa*. A revision of the structure of fruticulic B. *Phytochemistry* **1989**, *28*, 567–570. [\[CrossRef\]](#)
129. Erbario, M.; Ehrenfried, C.A.; Stefanello, M.É.A.; Dos Santos, É.P. Morphoanatomical and phytochemical studies of *Salvia lachnostachys* (Lamiaceae). *Microsc. Res. Tech.* **2012**, *75*, 1737–1744. [\[CrossRef\]](#)
130. Bisio, A.; Romussi, G.; Russo, E.; De Tommasi, N.; Mascolo, N.; Alfieri, A.; Bonito, M.C.; Cicala, C. Platelet antiaggregating activity and chemical constituents of *Salvia x jamensis* J. Compton. *Nat. Prod. Commun.* **2008**, *3*, 881–884. [\[CrossRef\]](#)
131. Schito, A.; Piatti, G.; Stauder, M.; Bisio, A.; Giacomelli, E.; Romussi, G.; Pruzzo, C. Effects of demethylfruticuline A and fruticuline A from *Salvia corrugata* Vahl. on biofilm production in vitro by multiresistant strains of *Staphylococcus aureus*, *Staphylococcus epidermidis* and *Enterococcus faecalis*. *Int. J. Antimicrob. Agents* **2011**, *37*, 129–134. [\[CrossRef\]](#)

132. Santos Oliveira, C.; Alvarez, C.J.; Pereira Cabral, M.R.; Sarragiotto, M.H.; Salvador, M.J.; Alves Stefanello, M.É. Three new diterpenoids from the leaves of *Salvia lachnostachys*. *Nat. Prod. Res.* **2022**, *36*, 5600–5605. [CrossRef]
133. Piccinelli, A.C.; Figueiredo de Santana Aquino, D.; Morato, P.N.; Kuraoka-Oliveira, Â.M.; Strapasson, R.L.B.; Dos Santos, E.P.; Stefanello, M.É.A.; Oliveira, R.J.; Kassuya, C.A.L. Anti-inflammatory and antihyperalgesic activities of ethanolic extract and fruticulin A from *Salvia lachnostachys* leaves in mice. *Evid. Based Complement. Altern. Med.* **2014**, *2014*, 835914. [CrossRef]
134. Giannoni, P.; Narcisi, R.; De Toter, D.; Romussi, G.; Quarto, R.; Bisio, A. The administration of demethyl fruticulin A from *Salvia corrugata* to mammalian cells lines induces “anoikis”, a special form of apoptosis. *Phytomedicine* **2010**, *17*, 449–456. [CrossRef]
135. Monticone, M.; Bisio, A.; Daga, A.; Giannoni, P.; Giaretti, W.; Maffei, M.; Pfeffer, U.; Romeo, F.; Quarto, R.; Romussi, G. Demethyl fruticulin A (SCO-1) causes apoptosis by inducing reactive oxygen species in mitochondria. *J. Cell. Biochem.* **2010**, *111*, 1149–1159. [CrossRef]
136. Hadavand Mirzaei, H.; Hosseini, S.M. In-silico assessments of fruticulin-a and demethylfruticulin-a isolated from *Salvia* species against important anticancer targets. *RJP* **2023**, *10*, 31–41.
137. Rhodes, M.J.C.; Parr, A.J.; Giulietti, A.; Aird, E.L.H. Influence of exogenous hormones on the growth and secondary metabolite formation in transformed root cultures. *Plant Cell Tissue Organ Cult.* **1994**, *38*, 143–151. [CrossRef]
138. Nussbaumer, P.; Kapetanidis, I.; Christen, P. Hairy roots of *Datura candida* × *D. aurea*: Effect of culture medium composition on growth and alkaloid biosynthesis. *Plant Cell Rep.* **1998**, *17*, 405–409. [CrossRef]
139. Poulev, A.; O’Neal, J.M.; Logendra, S.; Pouleva, R.B.; Timeva, V.; Garvey, A.S.; Gleba, D.; Jenkins, I.S.; Halpern, B.T.; Kneer, R. Elicitation, a new window into plant chemodiversity and phytochemical drug discovery. *J. Med. Chem.* **2003**, *46*, 2542–2547. [CrossRef]
140. Smetanska, I. Production of secondary metabolites using plant cell cultures. *Adv. Biochem. Eng./Biotechnol.* **2008**, *111*, 187–228. [CrossRef]
141. Akula, R.; Ravishankar, G.A. Influence of abiotic stress signals on secondary metabolites in plants. *Plant Signal. Behav.* **2011**, *6*, 1720–1731. [CrossRef]
142. Yue, W.; Ming, Q.-L.; Lin, B.; Rahman, K.; Zheng, C.-J.; Han, T.; Qin, L.-P. Medicinal plant cell suspension cultures: Pharmaceutical applications and high-yielding strategies for the desired secondary metabolites. *Crit. Rev. Biotechnol.* **2016**, *36*, 215–232. [CrossRef]
143. Murashige, T.; Skoog, F. A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol. Plant.* **1962**, *15*, 473–497. [CrossRef]
144. Gamborg, O.L.C.; Miller, R.A.; Ojima, K. Nutrient requirements of suspension cultures of soybean root cells. *Exp. Cell Res.* **1968**, *50*, 151–158. [CrossRef]
145. Veliky, I.A.; Martin, S.M. A fermenter for plant cell suspension cultures. *Can. J. Microbiol.* **1970**, *16*, 223–226. [CrossRef]
146. Efferth, T. Biotechnology Applications of Plant Callus Cultures. *Engineering* **2019**, *5*, 50–59. [CrossRef]
147. Guo, X.; Gao, W.; Li, K. Tissue culture of *Salvia miltiorrhiza* adventitious roots (II) Effects of carbon, nitrogen, and phosphate sources on culture of *Salvia miltiorrhiza* adventitious roots. *Chin. Tradit. Herb. Drugs* **1994**, wpr-576777. Available online: <https://www.kczg.org.cn/scholar/paperDetail?uid=115791&id=52132373> (accessed on 15 February 2024).
148. Gupta, S.K.; Liu, R.-B.; Liaw, S.-Y.; Chan, H.-S.; Tsay, H.-S. Enhanced tanshinone production in hairy roots of ‘*Salvia miltiorrhiza* Bunge’ under the influence of plant growth regulators in liquid culture. *Bot. Stud.* **2011**, *52*, 435–443.
149. Rao, S.R.; Ravishankar, G. Plant cell cultures: Chemical factories of secondary metabolites. *Biotechnol. Adv.* **2002**, *20*, 101–153.
150. Abbasi, B.H.; Tian, C.-L.; Murch, S.J.; Saxena, P.K.; Liu, C.-Z. Light-enhanced caffeic acid derivatives biosynthesis in hairy root cultures of *Echinacea purpurea*. *Plant Cell Rep.* **2007**, *26*, 1367–1372. [CrossRef]
151. Ali, M.; Abbasi, B.H. Light-induced fluctuations in biomass accumulation, secondary metabolites production and antioxidant activity in cell suspension cultures of *Artemisia absinthium* L. *J. Photochem. Photobiol. B Biol.* **2014**, *140*, 223–227. [CrossRef]
152. Liu, C.; Zhao, Y.; Wang, Y. Artemisinin: Current state and perspectives for biotechnological production of an antimalarial drug. *Appl. Microbiol. Biotechnol.* **2006**, *72*, 11–20. [CrossRef]
153. Yang, L.; Wen, K.-S.; Ruan, X.; Zhao, Y.-X.; Wei, F.; Wang, Q. Response of plant secondary metabolites to environmental factors. *Molecules* **2018**, *23*, 762. [CrossRef]
154. Tabata, M.; Mizukami, H.; Hiraoka, N.; Konoshima, M. Pigment formation in callus cultures of *Lithospermum erythrorhizon*. *Phytochemistry* **1974**, *13*, 927–932. [CrossRef]
155. Chandra, S.; Chandra, R. Engineering secondary metabolite production in hairy roots. *Phytochem. Rev.* **2011**, *10*, 371–395. [CrossRef]
156. Radman, R.; Saez, T.; Bucke, C.; Keshavarz, T. Elicitation of plants and microbial cell systems. *Biotechnol. Appl. Biochem.* **2003**, *37*, 91–102. [CrossRef]
157. Soundarapandian, S.; Dhandayuthapani, K. Interaction of plant growth promoting rhizobacteria (PGPR) and endophytes with medicinal plants-new avenues for phytochemicals. *J. Phytol.* **2010**, *2*, 91–100.
158. Nartop, P. Engineering of biomass accumulation and secondary metabolite production in plant cell and tissue cultures. In *Plant Metabolites and Regulation under Environmental Stress*; Ahmad, P., Ahanger, M.A., Singh, V.P., Tripathi, D.K., Alam, P., Alyemeni, M.N., Eds.; Elsevier: Amsterdam, The Netherlands, 2018; pp. 169–194.
159. Ganapathi, B.; Kargi, F. Recent advances in indole alkaloid production by *Catharanthus roseus* (Periwinkle). *J. Exp. Bot.* **1990**, *41*, 259–267. [CrossRef]

160. Naik, P.M.; Al-Khayri, J.M. Abiotic and biotic elicitors-role in secondary metabolites production through in vitro culture of medicinal plants. In *Abiotic and Biotic Stress in Plants—Recent Advances and Future Perspectives*; Shanker, A.K., Shanker, C., Eds.; InTech: Vienna, Austria, 2016; pp. 247–277.
161. Ferrari, S. Biological elicitors of plant secondary metabolites: Mode of action and use in the production of nutraceuticals. *Adv. Exp. Med. Biol.* **2010**, *698*, 152–166. [\[CrossRef\]](#)
162. Giauque, H.; Hawkes, C.V. Climate affects symbiotic fungal endophyte diversity and performance. *Am. J. Bot.* **2013**, *100*, 1435–1444. [\[CrossRef\]](#)
163. Tyc, O.; Song, C.; Dickschat, J.S.; Vos, M.; Garbeva, P. The ecological role of volatile and soluble secondary metabolites produced by soil bacteria. *Trends Microbiol.* **2017**, *25*, 280–292. [\[CrossRef\]](#)
164. Adegboye, M.F.; Babalola, O.O. Taxonomy and ecology of antibiotic producing actinomycetes. *Afr. J. Agric. Res.* **2012**, *7*, 2255–2261.
165. Olanrewaju, O.S.; Babalola, O.O. *Streptomyces*: Implications and interactions in plant growth promotion. *Appl. Microbiol. Biotechnol.* **2019**, *103*, 1179–1188. [\[CrossRef\]](#)
166. Shweta, S.; Gurumurthy, B.R.; Ravikanth, G.; Ramanan, U.S.; Shivanna, M.B. Endophytic fungi from *Miquelia dentata* Bedd., produce the anti-cancer alkaloid, camptothecine. *Phytomedicine* **2013**, *20*, 337–342. [\[CrossRef\]](#)
167. Zhao, J.; Xue, Q.-H.; Wang, L.-N.; Duan, C.-M.; Xue, L.; Mao, N. Antagonistic effect of multifunctional actinomycete strain Act12 on soil-borne pathogenic fungi and its identification. *Chin. J. Eco-Agric.* **2011**, *19*, 394–398. [\[CrossRef\]](#)
168. de Jesus Sousa, J.A.; Olivares, F.L. Plant growth promotion by streptomycetes: Ecophysiology, mechanisms and applications. *Chem. Biol. Technol. Agric.* **2016**, *3*, 24.
169. Yan, Y.; Zhang, S.; Yang, D.; Zhang, J.; Liang, Z. Effects of *Streptomyces pactum* Act12 on *Salvia miltiorrhiza* hairy root growth and tanshinone synthesis and its mechanisms. *Appl. Biochem. Biotechnol.* **2014**, *173*, 883–893. [\[CrossRef\]](#)
170. Barea, J.-M.; Pozo, M.J.; Azcon, R.; Azcon-Aguilar, C. Microbial co-operation in the rhizosphere. *J. Exp. Bot.* **2005**, *56*, 1761–1778. [\[CrossRef\]](#)
171. Paterson, J.; Jahanshah, G.; Li, Y.; Wang, Q.; Mehnaz, S.; Gross, H. The contribution of genome mining strategies to the understanding of active principles of PGPR strains. *FEMS Microbiol. Ecol.* **2016**, *93*, 249. [\[CrossRef\]](#)
172. Kloepper, J.W.; Ryu, C.-M.; Zhang, S. Induced systemic resistance and promotion of plant growth by *Bacillus* spp. *Phytopathology* **2004**, *94*, 1259–1266. [\[CrossRef\]](#) [\[PubMed\]](#)
173. Zhao, J.L.; Zhou, L.G.; Wu, J.Y. Promotion of *Salvia miltiorrhiza* hairy root growth and tanshinone production by polysaccharide-protein fractions of plant growth-promoting rhizobacterium *Bacillus cereus*. *Process Biochem.* **2010**, *45*, 1517–1522. [\[CrossRef\]](#)
174. Wu, J.-Y.; Ng, J.; Shi, M.; Wu, S.-J. Enhanced secondary metabolite (tanshinone) production of *Salvia miltiorrhiza* hairy roots in a novel root-bacteria coculture process. *Appl. Microbiol. Biotechnol.* **2007**, *77*, 543–550. [\[CrossRef\]](#)
175. Yan, Y.; Zhang, S.; Zhang, J.; Ma, P.; Duan, J.-L.; Liang, Z. Effect and mechanism of endophytic bacteria on growth and secondary metabolite synthesis in *Salvia miltiorrhiza* hairy roots. *Acta Physiol. Plant.* **2014**, *36*, 1095–1105. [\[CrossRef\]](#)
176. Zhai, X.; Luo, D.; Li, X.; Han, T.; Jia, M.; Kong, Z.; Ji, J.; Rahman, K.; Qin, L.; Zheng, C. Endophyte *Chaetomium globosum* D38 promotes bioactive constituents accumulation and root production in *Salvia miltiorrhiza*. *Front. Microbiol.* **2018**, *8*, 2694. [\[CrossRef\]](#)
177. Harman, G.E.; Howell, C.R.; Viterbo, A.; Chet, I.; Lorito, M. *Trichoderma* species—Opportunistic, avirulent plant symbionts. *Nat. Rev. Microbiol.* **2004**, *2*, 43. [\[CrossRef\]](#)
178. Ming, Q.; Han, T.; Li, W.; Zhang, Q.; Zhang, H.; Zheng, C.; Huang, F.; Rahman, K.; Qin, L. Tanshinone IIA and tanshinone I production by *Trichoderma atroviride* D16, an endophytic fungus in *Salvia miltiorrhiza*. *Phytomedicine* **2012**, *19*, 330–333. [\[CrossRef\]](#) [\[PubMed\]](#)
179. Ming, Q.; Su, C.; Zheng, C.; Jia, M.; Zhang, Q.; Zhang, H.; Rahman, K.; Han, T.; Qin, L. Elicitors from the endophytic fungus *Trichoderma atroviride* promote *Salvia miltiorrhiza* hairy root growth and tanshinone biosynthesis. *J. Exp. Bot.* **2013**, *64*, 5687–5694. [\[CrossRef\]](#)
180. Rabea, E.I.; Badawy, M.E.-T.; Stevens, C.V.; Smagghe, G.; Steurbaut, W. Chitosan as antimicrobial agent: Applications and mode of action. *Biomacromolecules* **2003**, *4*, 1457–1465. [\[CrossRef\]](#)
181. Zhao, J.-L.; Zhou, L.-G.; Wu, J.-Y. Effects of biotic and abiotic elicitors on cell growth and tanshinone accumulation in *Salvia miltiorrhiza* cell cultures. *Appl. Microbiol. Biotechnol.* **2010**, *87*, 137–144. [\[CrossRef\]](#) [\[PubMed\]](#)
182. Ramirez-Estrada, K.; Vidal-Limon, H.; Hidalgo, D.; Moyano, E.; Golenioswki, M.; Cusidó, R.; Palazon, J. Elicitation, an effective strategy for the biotechnological production of bioactive high-added value compounds in plant cell factories. *Molecules* **2016**, *21*, 182. [\[CrossRef\]](#)
183. Yan, J.; Li, H.; Li, S.; Yao, R.; Deng, H.; Xie, Q.; Xie, D. The Arabidopsis F-box protein CORONATINE INSENSITIVE1 is stabilized by SCFCOI1 and degraded via the 26S proteasome pathway. *Plant Cell* **2013**, *25*, 486–498. [\[CrossRef\]](#) [\[PubMed\]](#)
184. Weiler, E.; Kutchan, T.; Gorba, T.; Brodschelm, W.; Niesel, U.; Bublitz, F. The *Pseudomonas* phytotoxin coronatine mimics octadecanoid signalling molecules of higher plants. *FEBS Lett.* **1994**, *345*, 9–13. [\[CrossRef\]](#)
185. Vaccaro, M.C.; Mariaevelina, A.; Malafronte, N.; De Tommasi, N.; Leone, A. Increasing the synthesis of bioactive abietane diterpenes in *Salvia sclarea* hairy roots by elicited transcriptional reprogramming. *Plant Cell Rep.* **2017**, *36*, 375–386. [\[CrossRef\]](#)
186. Li, G.-J.; Wang, S.-C.; Xia, K.; Zhou, X. Effect of yeast elicitor and salicylic acid on the fluctuation of phytohormone contents in Ti-transformed *Salvia miltiorrhiza* cell cultures. *Plant Growth Regul.* **2003**, *39*, 27–32. [\[CrossRef\]](#)

187. Miyasaka, H.; Nasu, M.; Yoneda, K. *Salvia miltiorrhiza*: In vitro production of cryptotanshinone and ferruginol. In *Medicinal and Aromatic Plants II*; Springer: Berlin/Heidelberg, Germany, 1989; pp. 417–430.
188. Chen, H.; Chen, F. Effects of methyl jasmonate and salicylic acid on cell growth and cryptotanshinone formation in Ti transformed *Salvia miltiorrhiza* cell suspension cultures. *Biotechnol. Lett.* **1999**, *21*, 803–807. [\[CrossRef\]](#)
189. Shi, M.; Kwok, K.W.; Wu, J.Y. Enhancement of tanshinone production in *Salvia miltiorrhiza* Bunge (red or Chinese sage) hairy-root culture by hyperosmotic stress and yeast elicitor. *Biotechnol. Appl. Biochem.* **2007**, *46*, 191–196. [\[CrossRef\]](#)
190. Ge, X.; Wu, J. Induction and potentiation of diterpenoid tanshinone accumulation in *Salvia miltiorrhiza* hairy roots by β -aminobutyric acid. *Appl. Microbiol. Biotechnol.* **2005**, *68*, 183–188. [\[CrossRef\]](#)
191. Wu, J.-Y.; Shi, M. Ultrahigh diterpenoid tanshinone production through repeated osmotic stress and elicitor stimulation in fed-batch culture of *Salvia miltiorrhiza* hairy roots. *Appl. Microbiol. Biotechnol.* **2008**, *78*, 441–448. [\[CrossRef\]](#)
192. Yan, Q.; Hu, Z.; Tan, R.X.; Wu, J. Efficient production and recovery of diterpenoid tanshinones in *Salvia miltiorrhiza* hairy root cultures with in situ adsorption, elicitation and semi-continuous operation. *J. Biotechnol.* **2005**, *119*, 416–424. [\[CrossRef\]](#)
193. Yang, D.; Fang, Y.; Xia, P.; Zhang, X.; Liang, Z. Diverse responses of tanshinone biosynthesis to biotic and abiotic elicitors in hairy root cultures of *Salvia miltiorrhiza* and *Salvia castanea* Diels f. *tomentosa*. *Gene* **2018**, *643*, 61–67. [\[CrossRef\]](#)
194. Chen, H.; Chena, F.; Chiu, F.C.; Lo, C.M. The effect of yeast elicitor on the growth and secondary metabolism of hairy root cultures of *Salvia miltiorrhiza*. *Enzyme Microb. Technol.* **2001**, *28*, 100–105. [\[CrossRef\]](#)
195. Owolabi, I.O.; Yupanqui, C.T.; Siripongvutikorn, S. Enhancing secondary metabolites (emphasis on phenolics and antioxidants) in plants through elicitation and metabolomics. *Pak. J. Nutr.* **2018**, *17*, 411–420.
196. Bian, L.H.; Zou, L.; Zhou, B.Q.; Liu, W.; Zhou, J.; Wang, X. Effect of Lanthanum on accumulation of active constituent and key enzymes expression of *Salvia miltiorrhiza* hairy root. *Zhongguo Zhong Yao Za Zhi* **2016**, *41*, 4344–4349.
197. Han, M.; Guo, W.; Liang, Z.; Yang, D.; Yan, X.; Zhu, Y.; Liu, Y. Effects of cerous nitrate on growth and tanshinone production in *Salvia miltiorrhiza* hairy roots. *J. Rare Earths* **2015**, *33*, 1228–1235. [\[CrossRef\]](#)
198. Zhang, C.; Yan, Q.; Cheuk, W.-K.; Wu, J. Enhancement of tanshinone production in *Salvia miltiorrhiza* hairy root culture by Ag^+ elicitation and nutrient feeding. *Planta Medica* **2004**, *70*, 147–151.
199. Yu, Y.; Wang, T.; Wu, Y.; Zhou, Y.; Jiang, Y.; Zhang, L. Effect of elicitors on the metabolites in the suspension cell culture of *Salvia miltiorrhiza* Bunge. *Physiol. Mol. Biol. Plants* **2019**, *25*, 229–242. [\[CrossRef\]](#)
200. Rohwer, C.L.; Erwin, J.E. Horticultural applications of jasmonates. *J. Hortic. Sci. Biotechnol.* **2008**, *83*, 283–304. [\[CrossRef\]](#)
201. Baenas, N.; García-Viguera, C.; Moreno, D.A. Elicitation: A tool for enriching the bioactive composition of foods. *Molecules* **2014**, *19*, 13541–13563. [\[CrossRef\]](#)
202. De Geyter, N.; Gholami, A.; Goormachtig, S.; Goossens, A. Transcriptional machineries in jasmonate-elicited plant secondary metabolism. *Trends Plant Sci.* **2012**, *17*, 349–359. [\[CrossRef\]](#)
203. Liang, Z.S.; Yang, D.F.; Liang, X.; Zhang, Y.J.; Liu, Y.; Liu, F.H. Roles of reactive oxygen species in methyl jasmonate and nitric oxide-induced tanshinone production in *Salvia miltiorrhiza* hairy roots. *Plant Cell Rep.* **2012**, *31*, 873–883. [\[CrossRef\]](#)
204. Hao, X.; Shi, M.; Cui, L.; Xu, C.; Zhang, Y.; Kai, G. Effects of methyl jasmonate and salicylic acid on tanshinone production and biosynthetic gene expression in transgenic *Salvia miltiorrhiza* hairy roots. *Biotechnol. Appl. Biochem.* **2015**, *62*, 24–31. [\[CrossRef\]](#)
205. Xing, B.; Liang, L.; Liu, L.; Hou, Z.; Yang, D.; Yan, K.; Zhang, X.; Liang, Z. Overexpression of SmbHLH148 induced biosynthesis of tanshinones as well as phenolic acids in *Salvia miltiorrhiza* hairy roots. *Plant Cell Rep.* **2018**, *37*, 1681–1692. [\[CrossRef\]](#)
206. Wang, X.Y.; Cui, G.H.; Huang, L.Q.; Qiu, D.Y. Effects of methyl jasmonate on accumulation and release of tanshinones in suspension cultures of *Salvia miltiorrhiza* hairy root. *Zhongguo Zhong Yao Za Zhi* **2007**, *32*, 300–302.
207. Kuźma, Ł.; Bruchajzer, E.; Wysockińska, H. Methyl jasmonate effect on diterpenoid accumulation in *Salvia sclarea* hairy root culture in shake flasks and sprinkle bioreactor. *Enzym. Microb. Technol.* **2009**, *44*, 406–410. [\[CrossRef\]](#)
208. Kračun-Kolarević, M.; Dmitrović, S.; Filipović, B.; Perić, M.; Mišić, D.; Simonović, A.; Todorović, S. Influence of sodium salicylate on rosmarinic acid, carnosol and carnosic acid accumulation by *Salvia officinalis* L. shoots grown in vitro. *Biotechnol. Lett.* **2015**, *37*, 1693–1701. [\[CrossRef\]](#)
209. Yang, D.; Sheng, D.; Duan, Q.; Liang, X.; Liang, Z.; Liu, Y. PEG and ABA trigger the burst of reactive oxygen species to increase tanshinone production in *Salvia miltiorrhiza* hairy roots. *J. Plant Growth Regul.* **2012**, *31*, 579–587. [\[CrossRef\]](#)
210. Zhou, J.; Ran, Z.; Xu, Z.; Liu, Q.; Huang, M.; Fang, L.; Guo, L. Effects of smoke-water and smoke-isolated karrikinolide on tanshinones production in *Salvia miltiorrhiza* hairy roots. *S. Afr. J. Bot.* **2018**, *119*, 265–270. [\[CrossRef\]](#)
211. Wang, C.H.; Zheng, L.P.; Tian, H.; Wang, J.W. Synergistic effects of ultraviolet-B and methyl jasmonate on tanshinone biosynthesis in *Salvia miltiorrhiza* hairy roots. *J. Photochem. Photobiol. B* **2016**, *159*, 93–100. [\[CrossRef\]](#)
212. Chen, I.J.; Lee, M.S.; Lin, M.K.; Ko, C.Y.; Chang, W.T. Blue light decreases tanshinone IIA content in *Salvia miltiorrhiza* hairy roots via genes regulation. *J. Photochem. Photobiol. B* **2018**, *183*, 164–171. [\[CrossRef\]](#)
213. Cheng, Q.; He, Y.; Li, G.; Liu, Y.; Gao, W.; Huang, L. Effects of combined elicitors on tanshinone metabolic profiling and SmCPS expression in *Salvia miltiorrhiza* hairy root cultures. *Molecules* **2013**, *18*, 7473–7485. [\[CrossRef\]](#)
214. Kai, G.; Liao, P.; Zhang, T.; Zhou, W.; Wang, J.; Xu, H.; Liu, Y.; Zhang, L. Characterization, expression profiling, and functional identification of a gene encoding geranylgeranyl diphosphate synthase from *Salvia miltiorrhiza*. *Biotechnol. Bioprocess Eng.* **2010**, *15*, 236–245. [\[CrossRef\]](#)
215. Hong, M.L.K.; Bhatt, A.; Ping, N.S.; Keng, C.L. Detection of elicitation effect on *Hyoscyamus niger* L. root cultures for the root growth and production of tropane alkaloids. *Rom. Biotechnol. Lett.* **2012**, *17*, 7340–7351.

216. Ge, X.; Wu, J. Tanshinone production and isoprenoid pathways in *Salvia miltiorrhiza* hairy roots induced by Ag⁺ and yeast elicitor. *Plant Sci.* **2005**, *168*, 487–491. [\[CrossRef\]](#)
217. Zhao, J.-L.; Zou, L.; Zhang, C.-Q.; Li, Y.-Y.; Peng, L.-X.; Xiang, D.-B.; Zhao, G. Efficient production of flavonoids in *Fagopyrum tataricum* hairy root cultures with yeast polysaccharide elicitation and medium renewal process. *Pharmacogn. Mag.* **2014**, *10*, 234. [\[CrossRef\]](#)
218. Wang, C.; Wu, J.; Mei, X. Enhancement of taxol production and excretion in *Taxus chinensis* cell culture by fungal elicitation and medium renewal. *Appl. Microbiol. Biotechnol.* **2001**, *55*, 404–410. [\[CrossRef\]](#)
219. Diwan, R.; Malpathak, N. Bioprocess optimization of furanocoumarin elicitation by medium renewal and re-elicitation: A perfusion-based approach. *Appl. Biochem. Biotechnol.* **2011**, *163*, 756–764. [\[CrossRef\]](#)
220. Chiang, L.; Abdullah, M.A. Enhanced anthraquinones production from adsorbent-treated *Morinda elliptica* cell suspension cultures in production medium strategy. *Process Biochem.* **2007**, *42*, 757–763. [\[CrossRef\]](#)
221. Yan, Q.; Wu, J.Y.; Liu, R. Modeling of tanshinone synthesis and phase distribution under the combined effect of elicitation and in situ adsorption in *Salvia miltiorrhiza* hairy root cultures. *Biotechnol. Lett.* **2011**, *33*, 813–819. [\[CrossRef\]](#)
222. Miyasaka, H.; Nasu, M.; Yamamoto, T.; Endo, Y.; Yoneda, K. Production of cryptotanshinone and ferruginol by immobilized cultured cells of *Salvia miltiorrhiza*. *Phytochemistry* **1986**, *25*, 1621–1624. [\[CrossRef\]](#)
223. Vuong, T.V.; Franco, C.; Zhang, W. Treatment strategies for high resveratrol induction in *Vitis vinifera* L. cell suspension culture. *Biotechnol. Rep.* **2014**, *1*, 15–21. [\[CrossRef\]](#)
224. Wei, T.; Gao, Y.; Deng, K.; Zhang, L.; Yang, M.; Liu, X.; Qi, C.; Wang, C.; Song, W.; Zhang, Y. Enhancement of tanshinone production in *Salvia miltiorrhiza* hairy root cultures by metabolic engineering. *Plant Methods* **2019**, *15*, 53. [\[CrossRef\]](#)
225. Cao, W.; Wang, Y.; Shi, M.; Hao, X.; Zhao, W.; Wang, Y.; Ren, J.; Kai, G. Transcription factor SmWRKY1 positively promotes the biosynthesis of tanshinones in *Salvia miltiorrhiza*. *Front. Plant Sci.* **2018**, *9*, 554. [\[CrossRef\]](#)
226. Zhang, J.; Zhou, L.; Zheng, X.; Zhang, J.; Yang, L.; Tan, R.; Zhao, S. Overexpression of SmMYB9b enhances tanshinone concentration in *Salvia miltiorrhiza* hairy roots. *Plant Cell Rep.* **2017**, *36*, 1297–1309. [\[CrossRef\]](#)
227. Shi, M.; Luo, X.; Ju, G.; Li, L.; Huang, S.; Zhang, T.; Wang, H.; Kai, G. Enhanced diterpene tanshinone accumulation and bioactivity of transgenic *Salvia miltiorrhiza* hairy roots by pathway engineering. *J. Agric. Food Chem.* **2016**, *64*, 2523–2530. [\[CrossRef\]](#)
228. Gu, X.-C.; Chen, J.-F.; Xiao, Y.; Di, P.; Xuan, H.-J.; Zhou, X.; Zhang, L.; Chen, W.-S. Overexpression of allene oxide cyclase promoted tanshinone/phenolic acid production in *Salvia miltiorrhiza*. *Plant Cell Rep.* **2012**, *31*, 2247–2259. [\[CrossRef\]](#)
229. Bai, Z.; Li, W.; Jia, Y.; Yue, Z.; Jiao, J.; Huang, W.; Xia, P.; Liang, Z. The ethylene response factor SmERF6 co-regulates the transcription of SmCPS1 and SmKSL1 and is involved in tanshinone biosynthesis in *Salvia miltiorrhiza* hairy roots. *Planta* **2018**, *248*, 243–255. [\[CrossRef\]](#)
230. Alfieri, M.; Vaccaro, M.C.; Cappetta, E.; Ambrosone, A.; De Tommasi, N.; Leone, A. Coactivation of MEP-biosynthetic genes and accumulation of abietane diterpenes in *Salvia sclarea* by heterologous expression of WRKY and MYC2 transcription factors. *Sci. Rep.* **2018**, *8*, 11009. [\[CrossRef\]](#)
231. Vaccaro, M.; Bernal, V.O.; Malafronte, N.; De Tommasi, N.; Leone, A. High yield of bioactive abietane diterpenes in *Salvia sclarea* hairy roots by overexpressing cyanobacterial DXS or DXR genes. *Planta Medica* **2019**, *85*, 973–980. [\[CrossRef\]](#)

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.